

Biologically-Induced Precipitation of Minerals in a Medium with Zinc Under Sulfate-Reducing Conditions

DOROTA WOLICKA¹, ANDRZEJ BORKOWSKI¹, URSZULA JANKIEWICZ², WOJCIECH STĘPIEŃ³
and PAWEŁ KOWALCZYK^{4,5*}

¹Institute of Geochemistry, Mineralogy and Petrology, Faculty of Geology, University of Warsaw, Warsaw, Poland

²Department of Biochemistry, Faculty of Agriculture and Biology, Warsaw University of Life Sciences – SGGW, Warsaw, Poland; ³Department of Soil Sciences, Faculty of Agriculture and Biology, Warsaw University of Life Sciences – SGGW, Warsaw, Poland; ⁴Department of Microbial Biology, Faculty of Agriculture and Biology, Warsaw University of Life Sciences – SGGW, Warsaw, Poland; ⁵Bionicum LTD, Warsaw, Poland

Submitted 14 November 2013, revised 6 September 2014, accepted 27 February 2015

Abstract

Sulfate-reducing microbial communities were enriched from soils collected in areas with crude-oil exploitation. Cultures were grown in modified Postgate C medium and minimal medium, with ethanol or lactate as an electron donor. The batch cultures were grown with addition of zinc in concentrations of 100–700 mg/l. A lack of increased protein concentration in the solutions compared with the control batch, was noted in cultures containing over 200 mg Zn²⁺/l. The 16S rRNA method was applied to determine the specific composition of the selected microorganism communities. The analysis indicated the presence of *Desulfovibrio* spp., *Desulfobulbus* spp. and *Desulfotomaculum* spp. in the communities. Diffractometric analysis indicated the presence of biogenic sphalerite in cultures with 100 and 200 mg Zn²⁺/l and elemental sulfur in cultures with 200 mg Zn²⁺/l. Other post culture sediments (300–700 mg Zn²⁺/l) contained only hopeite [Zn₃(PO₄)₂·4H₂O] formed abiotically during the experiment, which was confirmed by studies of the activity of sulfate-reducing microbial communities.

Key words: biogenic precipitation, biomineralization, 16S rRNA, sulfate-reducing bacteria, zinc

Introduction

In the topmost part of the lithosphere, zinc is a trace element that poses a serious hazard to the environment. Heavy metals, including zinc, migrate in the hypergenic zone during various petrogenetic and geochemical processes, resulting in their concentration but equally often in their dispersion. The environmental issue caused by the migration of heavy metals, including zinc, is complex. On the one hand, such migration causes disappearance of the metals from ore beds, but on the other hand, the metals appear in increasing concentrations in the exploitation area. Ore exploitation and further treatment processes linked with the utilization of the raw ore deposit result in the formation of post-exploitation and technological wastes containing significant amounts of metals. Industrial wastes containing zinc are generally formed during the production of batteries, paints, plastics, polymer stabilizers as well as in printing enterprises (Fosmire, 1990). Environments contaminated by zinc may be harmful. Effluents that are naturally generated in waste dumpsites often discharge to surface water reservoirs, watercourses and soil, caus-

ing potential hazard to the environment. There are also areas with concentrations of selected metals, e.g., in areas of crude-oil exploitation. At present, there is an urgent need to apply processes that will allow recycling of heavy metals from wastes and poor ores and at the same time will minimize their negative influence on living organisms. Working out of an effective method of zinc recycling, e.g., from spoil tips, requires knowledge of the geochemical and mineral-forming processes taking place in the environment that will allow determination of the stability of the resulting mineral phases. Densification of heavy metals and their forms in soils depends on numerous factors of the physical and chemical environment, e.g., the magnitude of adsorption, the presence of humic acids and other soil components, the pH, the redox potential and others. Additionally, studies should include the role of microorganisms, especially sulfate-reducing bacteria (SRB), in the formation of secondary metal sulfides. SRB are a diverse group of anaerobic microorganisms that have the ability to reduce oxidized sulfur compounds and to oxidize organic compounds (Postgate, 1984; Hao *et al.*, 1996). They are considered to be the main producers of

* Corresponding author: P. Kowalczyk, Bionicum LTD, Warsaw, Poland; e-mail: pawel.kowalczyk@bionicum.com.pl

hydrogen sulfide in the biosphere. Sulfides that are produced during dissimilatory sulfate reduction may react with metals, including zinc, creating various secondary sulfides. In recent years, SRB have been applied to neutralize acid mine water, which may contain heavy-metal cations; and sparingly soluble metal sulfides are formed under these conditions (Jonson and Halleberg, 2003; Luptakova and Kusnierova, 2005; Kaksonen and Puhakka, 2007; Ong *et al.*, 2010). Application of SRB to acid mine drainage may remove heavy metals such as zinc from this environment. A few authors have asked why different kinds of minerals are formed during biological processes, and when and what influences these processes. Knowledge of biogeochemical interactions in the environment allows us to describe the processes and to predict them. It was long considered that the high toxicity of many heavy metals affects microbiological precipitation of sulfides by SRB. This opinion changed in 1961 when Becking and Moore (1961) carried out an experiment in which salts of selected metals were added to a medium with a composition close to that of marine water. The results explicitly pointed to SRB participation in the formation of secondary mineral phases because the post-culture sediments contained sphalerite (ZnS), a product of SO_4^{2-} and ZnCO_3 biotransformation. Taking into account the possible geochemical reactions, it can be assumed that in sedimentary settings, heavy-metal ions are largely adsorbed by clay minerals and form metal-organic compounds. All these processes lead to decrease of the toxicity level of a given metal by decreasing its concentration in the solution, although the metal is still capable of reactions with hydrogen sulfide and sulfide formation. On the other hand, the products of biochemical processes may include extracellular polymeric substances (EPS), which are mixtures of polysaccharides, mucopolysaccharides and proteins produced by microorganisms. The composition of EPS produced by SRB may be modified by the presence of different forms of organic matter in the environment, which may influence the increase of the metal-binding capacity (Zinkevich *et al.*, 1996). So far, the microbiological, geochemical and mineralogical processes leading to the formation of ZnS in environments impacted by human activities remain unrecognized. Spherical aggregates of sphalerite are commonly observed in biofilm structures where anaerobic conditions prevail. These zones are usually dominated by SRB that are relatively tolerant to oxygen, representing the families *Desulfobacteraceae* and *Desulfovibrionaceae* (Labrenz *et al.*, 2000; Vainshtein *et al.*, 1992). The precipitation of sphalerite at low temperatures may take place in mine-water environments (Ledin and Pedersen, 1996; Drury, 1999; Moreau *et al.*, 2004). In order to determine the effect of zinc concentration on the activity of selected groups of microorganisms, experiments were

performed in which zinc chloride was added at various concentrations to SRB cultures. There are only a few reports that describe both the influence of SRB on the formation of mineral phases that contain zinc and its effect on SRB activity.

The present study focused on the role of SRB isolated from soils with increased zinc concentrations, collected in areas of crude-oil exploitation, in the formation of zinc sulfide at high concentrations of the metal. It also focused on the toxicity of zinc in relation to sulfate-reducing microbial communities in batch cultures containing easily accessible carbon sources for SRB, *i.e.*, lactate or ethanol.

Experimental

Materials and Methods

Selection and isolation of sulfate-reducing microbial communities. The microorganisms were enriched from soil polluted by crude oil and oil-derived products from SE Poland. SRB are commonly found in soils contaminated by crude oil due to the ability to metabolize the oil-derived products (Feio *et al.*, 2004). In tested soil samples, C total was 3.2%, total S was about 120 mg/kg dry weight, total N was about 700 mg/kg, and Zn^{2+} was 92 mg/kg. In the tested soil samples the Zn^{2+} ions were only determined. First, an Easycult S test (Orion Diagnostica Espoo of Finland) was made to check for the presence of sulfidogenic microorganism communities; next, the microorganisms were selected using the microcosm method. Soil samples (10 g) were inserted in 100-ml flasks and covered with 80 ml of the particular medium. Two types of media were applied: a modified Postgate C medium (without yeast extract and sodium citrate) and a minimal medium with lactate or ethanol as electron donors. The flasks were tightly closed and incubated in darkness for 6 weeks at room temperature (about 22°C) in order to select anaerobic, sulfidogenic microorganism communities capable of simultaneous biodegradation of the applied carbon sources and sulfate reduction. The obtained SRB community was the inoculum to the SRB cultures in main experiment.

Cultures of sulfate-reducing microbial communities. Anaerobic batch cultures in modified liquid Postgate C medium were carried out in 0.5 l glass bottles filled to 0.25 l volume. The bottles were tightly sealed with rubber stoppers pierced with needles connected permanently to syringes, which were used to introduce the inoculum and to collect samples under N_2 . The inoculum-to-medium ratio was 1:10. The anaerobic conditions in the cultures were controlled by addition of resazurin as the oxygen-level indicator. Violet

Table I
Experimental setup

Source of carbon	SRB cultures		Biotic control		Abiotic control (for diffractometric analysis)	
	Ethanol	Lactate	Ethanol	Lactate	Ethanol	Lactate
Zn ²⁺ concentration [mg/l]	100	100	0	0	100	100
	200	200			200	200
	500	500			500	500
	700	700			700	700
SRB inoculum	10%	10%	10%	10%	without inoculum	without inoculum
Repetitions	2 ×	2 ×	2 ×	2 ×	1 ×	1 ×

colour indicated that the culture contained oxygen, and its absence pointed to anaerobic conditions. The experiment was carried out in two variants, one with ethanol and the other with lactate as the sole carbon source (4000 mg/l). The control batch consisted of SRB cultures without zinc (biotic control) and cultures with zinc and without SRB (abiotic control). The abiotic controls were conducted in order to analyze of mineral phases formed without microbial activity. All the SRB cultures were stationary and were conducted in the modified Postgate C medium. The experiment and the chemical determinations in cultures were made in duplicate. The experimental setup is shown in Table I.

Media. A modified liquid Postgate C medium (Postgate, 1984), composed of: KH₂PO₄ (500 mg/l), NH₄Cl (1000 mg/l), CaCl₂ (60 mg/l), MgSO₄ (60 mg/l), FeSO₄ (100 mg/l), Na₂SO₄ (4500 mg/l) without yeast extract and citrate, and a minimal medium (Wolicka and Kowalski, 2006), composed of NH₄Cl (1000 mg/l) and Na₂SO₄ (4500 mg/l), were used in the experiment. Lactate (4000 mg/l) or ethanol (4000 mg/l) were added to both media as the sole carbon sources. Resazurin (1 mg/l) was added to all cultures in order to control the level of oxidation. The medium did not contain yeast extract or sodium citrate. Zn²⁺ was added in concentrations of 100, 200, 500 and 700 mg/l to a modified Postgate C medium. Zn²⁺ was added as zinc (II) chloride.

Sulfate determinations were made using the turbidimetric method after reaction with barium chloride in a Thermo spectrophotometer at $\lambda = 400$ nm wavelength (Greenberg *et al.*, 1985).

Protein determinations in the cultures were made using the Lowry method after a biuret test enhanced by the Folin-Ciocalteu reagent in a Thermo spectrophotometer at $\lambda = 670$ nm (Genesys 10Vis, Thermo). The samples (5 ml) for determination were previously sonicated (30 kHz, 30 s) in order to determine total protein from cultures. The measurement was performed as follows. The 1 ml of sonicated sample was put into glass tubes and 5 ml of reagent (49 ml 2% Na₂CO₃ in 0.1 M NaOH + 0.5 ml 2% potassium sodium tartrate

+ 0.5 ml 1% CuSO₄) was added. After 5 min., 0.2 ml of Folin-Ciocalteu reagent (POCH, Gliwice, Poland) was added and immediately mixed. After 5 min, 0.2 ml 6 M NaOH was added and mixed, then the absorbance of the colored solution was immediately measured spectrophotometrically. Protein measurements were used as an indicator of biomass.

Zinc determinations in stationary cultures. The determinations were made using the Thermo spectrophotometer with application of available kits for determining zinc concentrations (Merck Zinc Test with a pyridylazo naphthol derivative).

Analysis of the post-culture sediments. After incubation, the cultures were centrifuged at 10 000 × g, and the obtained post-culture sediment was dried at 30°C under N₂. The samples were next ground in an agate mortar, and their mineral composition was determined using X-ray powder diffraction in a diffractometer (Panalytical X' Pert PRO MPD). The diffractometric analysis was conducted on 10 samples of post-culture sediments; five were taken from cultures that used ethanol as the sole carbon source, and five were taken from cultures with lactate as the sole carbon source. Additionally, the analysis of sediments from abiotic control with Zn²⁺ (500 and 700 mg/l) were also conducted.

Molecular analysis of the selected sulfidogenous microorganism communities. The taxonomic composition of the sulfate-reducing bacterial communities was obtained using molecular analysis. Isolation of chromosome DNA and analysis of gene 16S rRNA fragments were carried out according to commonly applied procedures of Collins *et al.* (1991). Bacterial DNA was isolated from a fluid culture of microorganisms with a commercial kit for chromosomal DNA isolation (A&A Biotechnology). The purity and concentration of the resulting DNA preparation were determined spectrophotometrically at 260 nm. Primers specific for bacterial 16S rRNA (27F 5'-AGAGTTTGATCCTG-GCTCAG-3' and 1492R 5'-GGTTACCTTGTTAC-GACTT-3') were used to amplify a 1540-bp segment from the 16S rRNA gene. The PCR reaction was made

using the GeneAmp PCR reagent kit with AmpliTaq DNA polymerase (Invitrogen). Amplification products were purified using the Wizard Purification System (Promega) and analyzed by electrophoresis. After amplification, the material was sequenced using the ABI 3730 Genetic Analyzer with application of the Perkin Elmer sequencing kit. The resulting nucleotide sequences were compared with gene 16S rRNA sequences available in the National Centre for Biotechnology Information (NCBI) database using NCBI's Blast 2.0 program and showed 99% homology with the corresponding sequences among different anaerobic species.

Results and Discussion

The influence of zinc concentrations on the activity of selected SRB communities is presented in Figs. 1 and 2 and the changes of concentration of sulphate and protein in control cultures are presented in Fig. 3. The initial concentration of protein at the level of about 2000 mg/l was derived from the inoculum. A slight increase of protein concentration and decrease of sulfate concentration was observed in cultures where the Zn concentration was 100 mg/l on a medium with ethanol as the sole carbon source (Fig. 1). Such trends were not observed in the remaining cultures, regardless of the applied carbon source. A slight increase of protein concentration and decrease of sulfate concentration was noted in the control batch. Based on the obtained results, it can be stated, that the zinc concentration tolerated by the isolated SRB community was 100–200 mg/l.

The results may partially confirm existing literature data. Zinc is a metal that may hamper in higher concentrations the metabolic activity of various microorganisms, including SRB (Utgikar *et al.*, 2002), but on the other hand many mechanisms responsible for metal-ion resistance in bacteria have been described (Brocklehurst and Morby, 2000). SRB are effective in reducing the sulfate concentration and neutralizing its acidity. Furthermore, most of the heavy metals present in acidic mine drainage can be precipitated as insoluble sulfides using biogenic sulfide produced by sulfate reduction (Barton and Tomei, 1995; Costa *et al.*, 2008; Martins *et al.*, 2009a). The reported toxic concentrations of heavy metals to sulfate reducers range from a few mg/l to 100 mg/l (Loka Bharathi *et al.*, 1990; Poulson *et al.*, 1997; Utgikar *et al.*, 2001). Data in the literature indicate that various concentrations of zinc inhibit SRB activity. Radhika *et al.* (2006) estimated that the concentration of zinc lethal to SRB is about 210 mg/l. Castillo *et al.* (2012) isolated communities from sediments in two acid streams draining the Iberian Pyrite Belt, in which zinc occurred at con-




Fig. 1. Concentration changes of sulfate, protein and zinc in sulfidogenic microbial cultures with ethanol as the sole organic carbon source at variable initial concentrations of Zn²⁺ (100, 200, 500, 700 mg/l). Standard deviation has been marked




Fig. 2. Concentration changes of sulfate, protein and zinc in sulfidogenic microbial cultures with lactate as the sole organic carbon source at variable initial concentrations of Zn^{2+} (100, 200, 500, 700 mg/l). Standard deviation has been marked




Fig. 3. Concentration changes of sulfate and protein in control batches (without addition of Zn^{2+}) of sulfidogenic microbial cultures with ethanol (above) or lactate (below) as the sole organic carbon source. Standard deviation has been marked

centrations of 400 and 30 mg/l. According to Azabou *et al.* (2007), a zinc concentration of 400 mg/l is toxic for SRB and inhibits their activity; SRB can carry out metabolic processes at concentrations of up to 150 mg/l (Martins *et al.*, 2009b). Zinc inhibits electron transport in the respiration cycle of microorganisms, and its toxicity in comparison to such metals as Hg, Cd, Cu, Ni, Co and Pb is rather low. There is data in the literature on the influence of zinc on SRB activity, but there are no reports that simultaneously discuss the influence of zinc concentration on SRB activity and the types of the resulting mineral phases.

In the present study, the determination of the influence of Zn^{2+} on the mineral composition of post-culture sediments was conducted. The presence of zinc sulfide was determined in the post-culture sediments from cultures with ethanol as the sole carbon source and 100 mg Zn^{2+} /l (Fig. 4). In the culture, where the zinc concentration was 200 mg/l, sphalerite (ZnS) as well as elemental sulfur were observed. It could be that elemental sulfur is formed by sulfidogenic bacterial communities. In natural ecosystems, the sulfur cycle should be in balance, meaning that the amount of sulfide that is oxidized should correspond to the amount of sulfate that is reduced. Such a balance can be found in a sulfuretum. This is a syntrophical bacterial community in which H_2S produced by sulfate-reducing bacteria is reoxidized




Fig. 4. X-ray powder diffractograms of post-culture sediments in cultures of selected SRB communities on a modified Postgate medium with ethanol as the sole carbon source and with addition of Zn^{2+} in concentrations of 100–700 mg/l. Symbols: Zn, sphalerite ZnS ; Hp, hopeite $Zn_3(PO_4)_2 \cdot H_2O$; S, sulfur

by the sulfur-compound-oxidizing bacteria. This process is not as common in the natural environment as dissimilation sulfate reduction, but it should not be excluded (Roy and Trudinger, 1970; Hedderich *et al.*, 1999). The remaining cultures containing more than 200 mg Zn^{2+}/l contained hopeite [$Zn_3(PO_4)_2 \cdot H_2O$], which was probably formed by abiotic processes because the medium used for SRB growth contained phosphates (KH_2PO_4). The abiotic control batch did not contain any mineral phases except hopeite. Hopeite was detected in abiotic controls containing 500 and 700 mg Zn^{2+}/l .

On a medium with lactate as the sole carbon source (Fig. 2), significant SRB activity could not be observed, but a slight increase of protein content in the cultures worth noting, pointing to the development of microorganisms accompanying SRB that were capable of activity in the presence of high zinc concentrations. The significant decrease of zinc concentration may have been the effect of this microflora activity, although this fact cannot be unambiguously confirmed. The effect of inoculum can be appear only at initial stage of experiment, when it can be observed the slight decrease of zinc concentration. A considerable impediment in the analysis of the results is also the lack of determination of the minerals in post-culture sediments that could not be obtained using X-ray diffraction. This lack probably was caused by the low number of mineral phases in the post-culture sediment and precipitation of amorphous or nanostructured chemical compounds, which often accompany EPS organic compounds. Moreau *et al.* (2007) suggest that formation of mineral aggregates is induced by extracellular metal-

binding polypeptides and proteins. Disordered morphological features of zinc sulfides have been described by Gramp *et al.* (2007), who tested for formation of Zn sulfides in cultures of SRB.

The 16S rRNA method was applied in order to determine the specific composition of the selected microorganism communities. The analysis indicated the presence of *Desulfovibrio* spp., *Desulfobulbus* spp. and *Desulfotomaculum* spp. in the communities. Some of these microorganisms are capable of sulfate reduction to elemental sulfur, which was confirmed by diffractometric analysis of the post-culture sediments. Moreover, removal of hydrogen sulfide from the environment by formation of sulfides results in a decrease of the reducing conditions that are indispensable for SRB activity (Labrenz *et al.*, 2000).

Conclusions. The results supplement the knowledge of the mineral-forming processes taking place in cultures with various concentrations of zinc and indicate a possibly significant participation of SRB in the processes taking place in the natural environment under hypergenic conditions. Moreover, they indicate the possible application of SRB in the treatment of acid mine drainage, but the high concentrations of metals potentially can limit the SRB activity. Cultures with zinc showed strong inhibition of the activity of selected sulfidogenic communities in cultures containing 200 mg Zn^{2+}/l . The post-culture sediments of cultures in a Postgate medium with ethanol as the sole carbon source contained sphalerite (cultures containing 100 mg Zn^{2+}/l) and elemental sulfur (cultures contain-

ing 200 mg Zn²⁺/l). In the remaining cultures, in which the content of zinc ions was much higher, the post-culture sediments contained zinc phosphate formed by abiotic processes. The results confirm the participation of SRB in mineral-forming processes in environments containing zinc.

Acknowledgments

We would like to thank the Polish Oil & Gas Company, especially Jozef Potera and Tadeusz Kozimor from the Sanok Branch and Zbigniew Ząbkiewicz from Warsaw. This work was supported by the Institute of Geochemistry, Mineralogy and Petrology under BST Grant No. IGMiP-28-2012. The authors thank Reviewers for their remarks and comments which have improved this article.

Literature

- Azabou S., T. Mechichi and S. Sayadi.** 2007. Zinc precipitation by heavy-metal tolerant sulfate-reducing bacteria enriched on phosphogypsum as a sulfate source. *Minerals Engineering* 20: 173–178.
- Barton L.L. and F.A. Tomei.** 1995. Characteristics and activities of sulfate reducing bacteria, pp. 1–32. In: L. L. Barton (eds). *Sulfate Reducing Bacteria*. Plenum Press, Chap 1. New York.
- Becking L.G.M.B. and D. Moore.** 1961. Biogenic sulfides. *Economic Geology* 56: 259–272.
- Brocklehurst K.R. and A.P. Morby.** 2000. Metal-ion tolerance in *Escherichia coli*: analysis of transcriptional profiles by gene-array technology. *Microbiology* 146: 2277–2282.
- Castillo J., R. Pérez-López, M.A. Caraballo, J.M. Nieto, M. Martins, M.C. Costa, M. Olias, J.C. Cerón and R. Tucoulou.** 2012. Biologically-induced precipitation of sphalerite-wurtzite nanoparticles by sulfate-reducing bacteria: Implications for acid mine drainage treatment. *Science of the Total Environment* 423: 176–184.
- Collins M.D., S. Wallbanks, D.J. Lane, J. Shah, R. Nietupski, J. Smida, M. Dorsch and E. Stackebrandt.** 1991. Phylogenetic analysis of the genus *Listeria* based on reverse transcriptase sequencing of 16S rRNA. *International Journal of Systematic and Evolutionary Microbiology* 41: 240–246.
- Costa M.C., M. Martins, C. Jesus and J.C. Duarte.** 2008. Treatment of acid mine drainage by sulphate-reducing bacteria using low cost matrices. *Water Air Soil Pollut.* 189: 149–162.
- Drury W.J.** 1999. Treatment of acid mine drainage with anaerobic solid-substrate reactors. *Water Environmental Research* 71: 1244–1250.
- Feio M.J., V. Zinkevic, I.B. Beech, E. Llobet-Brossa, P. Eaton, J. Schmitt and J. Guezennec.** 2004. *Desulfovibrio alaskensis* sp. nov., a sulphate reducing bacterium from a soured oil reservoir. *International Journal of Systematic and Evolutionary Microbiology* 54: 1747–1752.
- Fosmire G.J.** 1990. Zinc toxicity. *American Journal Clinical Nutrition* 51: 225–222.
- Gramp J.P., J.M. Bigham, K. Sasaki and O.H. Touvien.** 2007. Formation of Ni- and Zn-sulfides in cultures of Sulfate-Reducing Bacteria. *Geomicrobiology Journal* 24: 609–614.
- Greenberg A.E., R.R. Trussell and L.S. Clesceri.** 1985. Standard methods for the examination of water and wastewater, pp. 11–20. APHA-AWWA-WPCF, Washington, DC.
- Hao O.J, J.M. Chen, L. Huang and R.L. Buglass.** 1996. Sulfate-reducing bacteria. *Critical Reviews in Environmental Science and Technology* 26: 155–187.
- Hedderich R., O. Klimmek, A. Kroger, M. Dirmeier, M. Kelller and O. Stetter.** 1999. Anaerobic respiration with elemental sulfur and with disulfides. *FEMS Microbiology Reviews* 22: 353–381.
- Jonson D.B. and K.B. Halleberg.** 2003. The microbiology of acidic mine wastewaters. *Res. Microbiology* 154: 466–473.
- Kaksonen A.H. and J.A. Puhakka.** 2007. Sulfate reduction based bioprocesses for the treatment of acid mine drainage and the recovery of metals. *Engineering in Life Sciences* 7: 541–564.
- Labrenz M., G.K. Druschel, T. Thomsen-Ebert, B. Gilbert, S.A. Welch, K.M. Kemmer, G.A. Logan, R.E. Summons, G. De Stasio, P.L. Bond and others.** 2000. Formation of sphalerite (ZnS) deposits in natural biofilms of sulfate-reducing bacteria. *Science* 290: 1744–1747.
- Ledin M. and K. Pedersen.** 1996. The environmental impact of mine wastes – Roles of microorganisms and their significance in treatment of mine wastes. *Earth Science Reviews* 41: 67–108.
- Loka Bharathi P.A., V. Sathe and D. Chandramohan.** 1990. Effect of lead, mercury and cadmium on a sulphate-reducing bacterium. *Environmental Pollutants* 67 (4): 361–74.
- Luptakova A. and M. Kusnierova.** 2005. Bioremediation of acid mine drainage contaminated by SRB. *Hydrometallurgy* 77: 97–102.
- Martins M., M.L. Faleiro, R.J. Barros, A.R. Verissimo, M.A. Barreiros and M.C. Costa.** 2009a. Characterization and activity studies of highly heavy metal resistant sulphate-reducing bacteria to be used in acid mine drainage decontamination. *Journal of Hazardous Materials* 166: 706–713.
- Martins M., M.L. Faleiro, R.J. Barros, A.R. Verissimo and M.C. Costa.** 2009b. Biological sulphate reduction using food industry wastes as carbon sources. *Biodegradation Journal* 20 (4): 559–567.
- Moreau J.W., P.K. Weber, M.C. Martin, B. Gilbert, I.D. Hutcheon and J.F. Banfield.** 2007. Extracellular proteins limit the dispersal of biogenic nanoparticles. *Science* 316: 1600–1603.
- Moreau J.W., R.I. Webb and J.F. Banfield.** 2004. Ultrastructure, aggregation-state, and crystal growth of biogenic nanocrystalline sphalerite and wurtzite. *American Mineralogist* 89: 950–960.
- Ong S-A., E. Toorisaka, M. Hirata and T. Hano.** 2010. Adsorption and toxicity of heavy metals on activated sludge. *Science Asia* 36: 204–209.
- Postgate J.R.** 1984. The sulphate reducing bacteria, pp. 1–159. 2nd ed. *Cambridge University Press*, Cambridge.
- Poulson S.R., P.J.S. Colberg and J.I. Drever.** 1997. Toxicity of heavy metals (Ni, Zn) to *Desulfovibrio desulfuricans*. *Geomicrobiology Journal* 14: 41–49.
- Radhika V., S. Subramanian and K.A. Natarajan.** 2006. Bioremediation of zinc using *Desulfotomaculum nigrificans*: Bioprecipitation and characterization studies. *Water Research* 40: 3628–3636.
- Roy A.B and P.A. Trudinger.** 1970. The biochemistry of inorganic compounds of sulphur, pp. 1–399. *Cambridge University Press*, Cambridge.
- Utgikar V.P., B-Y. Chen, N. Chaudhary, H.H. Tabak, J.R. Hainjes and R. Govind.** 2001. Acute toxicity of heavy metals to acetate-utilizing mixed cultures of sulfate-reducing bacteria: EC100 and EC500. *Environmental Toxicology and Chemistry* 20: 2662–2669.
- Utgikar V.P., S.M. Harmon, N. Chaudhary, H.H. Tabak, R. Govind and J.R. Haines.** 2002. Inhibition of sulfate-reducing bacteria by metal sulfide formation in bioremediation of acid mine drainage. *Environmental Toxicology* 17: 40–48.
- Vainshtein M., H. Hippe and R.M. Kroppenstedt.** 1992. Cellular fatty acid composition of *Desulfovibrio* species and its use in classification of sulfate-reducing bacteria. *Syst. Applied Microbiology* 15: 554–556.
- Wolicka D. and W. Kowalski.** 2006. Biotransformation of phosphogypsum in petroleum-refining wastewaters. *Polish J. Environ. Stud.* 15 (2): 355–360.
- Zinkevich V., I. Bogdarina, H. Kang, M. Hill, R. Tapper and I.B. Beech.** 1996. Characterisation of exopolymers produced by different isolates of marine sulphate-reducing bacteria. *Int. Biodeterior Biodegrad* 37: 163–172.

