

## Safety Evaluation of Enterocin Producer *Enterococcus* sp. Strains Isolated from Traditional Turkish Cheeses

MINE AVCI and BANU ÖZDEN TUNCER\*

<sup>1</sup>Süleyman Demirel University, Faculty of Engineering, Department of Food Engineering, Isparta, Turkey

Submitted 26 July 2016, revised 14 October 2016, accepted 12 January 2017

### Abstract

The purpose of this study was to determine the antimicrobial activity and occurrence of bacteriocin structural genes in *Enterococcus* spp. isolated from different cheeses and also investigate some of their virulence factors. *Enterococcus* strains were isolated from 33 different cheeses. *Enterococcus faecium* (6 strains) and *Enterococcus faecalis* (5 strains) enterocin-producing strains were identified by 16S rDNA analyses. Structural genes *entA*, *entB*, *entP* and *entX* were detected in some isolates. Multiple enterocin structural genes were found in 7 strains. None of the tested enterococci demonstrated any  $\beta$ -haemolytic activity and only one strain had gelatinase activity. Six strains showed multiple antibiotic resistance patterns and in addition, *vanA* and several virulence genes were detected in many strains. Only *E. faecalis* MBE1-9 showed tyrosine decarboxylase activity and *tdc* gene was detected only in this strain.

**Key words:** *Enterococcus* sp. from cheeses, antibiotic resistance of food borne enterococci bacteriocin production ability, virulence factors

### Introduction

Enterococci are an important group of lactic acid bacteria (LAB) that inhabit the digestive tract of humans and animals as well as the surface of waters, soil and plants. They are also found in foods, especially of animal origin, such as traditional cheeses produced in different European and Mediterranean countries (Manolopoulou *et al.*, 2003; Yogurtcu and Tuncer, 2013). *Enterococcus faecalis*, *Enterococcus faecium* and *Enterococcus durans* are the most frequently isolated strains from traditional cheeses, and they play an important role for ripening of these cheeses based on their proteolysis, lipolysis, citrate breakdown abilities and hence they promote product's unique typical taste and flavor formation (Giraffa, 2002; 2003; Moreno *et al.*, 2006). They can also inhibit the growth of the food-borne pathogens and spoilage microorganisms by producing bacteriocin known as enterocins. The frequently encountered enterocins produced by enterococci strains are enterocins A, B, P, AS-48, L50A, L50B, 1071A, 1071B, and Q (Moreno *et al.*, 2006; Nes *et al.*, 2007; Edalatian *et al.*, 2012; Özden-Tuncer *et al.*, 2013).

Although enterococci have desirable technological and metabolic traits, in recent years they have been found to be associated with clinical infections such

as endocarditis, bacteremia and urinary tract. Therefore, enterococci have emerged as an important nosocomial pathogen in recent decades (Franz *et al.*, 2003; Moreno *et al.*, 2006; Peters *et al.*, 2007; Lindenstrauß *et al.*, 2011). Several putative virulence factors have been described in enterococci, such as gelatinase (*gelE*), cell wall adhesin (*efaA<sub>fm</sub>* and *efaA<sub>fs</sub>* from *E. faecium* and *E. faecalis*, respectively), sex pheromones (*cpd*, *cob*, *ccf*, and *cad*), collagen adhesin (*ace*), enterococcal surface protein (*esp<sub>fm</sub>* and *esp<sub>fs</sub>* from *E. faecium* and *E. faecalis*, respectively), aggregation substance (*agg*), and cytolyisin (*cylM*, *cylB*, and *cylA*) (Vankerckhoven *et al.*, 2004; Cariolato *et al.*, 2008; Chuang *et al.*, 2009; Barbosa *et al.*, 2010; Ben Belgacem *et al.*, 2010; Lindenstrauß *et al.*, 2011; Yogurtcu and Tuncer, 2013). In addition, enterococci are resistant to several antibiotics due to intrinsic and/or acquire genes located on the chromosomal DNA, plasmids or transposons. The interesting point about antibiotic resistance of enterococci is the increasing number of vancomycin and multiple antibiotic-resistant strains. The reason for the increase in multiple antibiotic resistance in enterococci is transfer of genetic elements carrying antibiotic resistance genes between pathogenic and non-pathogenic bacteria in the human or animal intestinal tract or even in food (Riboldi *et al.*, 2009; Sparo *et al.*, 2011).

\* Corresponding author: B. Özden Tuncer, Süleyman Demirel University, Faculty of Engineering, Department of Food Engineering, Isparta, Turkey; e-mail: banutuncer@sdu.edu.tr

Although the technological and metabolic activity of enterococci is essential for some cheeses, the presence of virulence factors and antibiotic resistance in enterococci should be investigated due to their ability of transferring virulence genes and antibiotic resistance determinants through the food chain. In recent years, virulence factors and antibiotic resistance of clinical isolates were mostly examined, but every day interest in food-borne enterococci and their pathogenicity is increasing. From these perspectives, the aim of this study was to isolate enterocin producing enterococci strains from various traditional artisanal cheeses in Turkey, and to determine their enterocin genes, antibiotic resistance patterns and virulence factors. Enterocin production, antibiotic resistance and virulence determinants of enterococci isolated from some cheese types in this study were investigated for the first time.

## Experimental

### Materials and Methods

**Cheese samples.** Cheese samples were obtained from various provinces (Denizli, Isparta, Balıkesir, Bolu, Kayseri, Van and Manisa) of Turkey and it has been noted that they are especially homemade. A total of 33 cheeses samples, including traditional Turkish white cheese, Tulum cheese, Manyas Kelle and Van Otlu (Herby) cheese made from cow's, sheep's and goat's milk were used in the present study.

**Isolation and phenotypically identification of *Enterococcus* sp. strains.** Ten grams of each cheese sample were homogenized in 90 ml sterile saline solution (0.85% NaCl, w/v) with waring blender (8011 ES HGB2WTS3, Torrington, Connecticut, 06790, USA). Serial decimal solutions were prepared in saline solution and spread onto Kanamycin Eascaline Azide Agar (Lab M, Ltd., Bury, Lancashire, U.K.) and incubated at 37°C for 48 h. Over an incubation period black colonies of presumptive *Enterococcus* sp. strains were randomly picked up and grown in de Man Rogosa Sharpe (MRS) broth medium (Lab M) at 37°C for 24 h. A total of 100 isolates were identified to the genus level by Gram staining, catalase and cultural tests such as growth in MRS broth (10°C, 37°C, 45°C and pH 9.6), tolerance of 6.5% (w/v) NaCl and resistance to heat (60°C for 30 and 60 min).

**Detection of bacteriocin production ability.** Antimicrobial activities of the isolates were evaluated towards selected indicator bacteria by using the toothpick method as described by van Belkum *et al.* (1989). Briefly, overnight presumptive *Enterococcus* sp. strains were grown on MRS solid medium at 37°C for 18 h. After incubation time, sterile toothpicks were used to

ensure create spots on MRS agar. After the incubation period, an indicator lawn of 5 ml of appropriate soft agar (0.5%, w/v), containing 100 µl indicator bacteria was poured onto the surface and incubated at the required temperature for indicators for 18 h. After incubation, all inhibition zones around the colonies were examined and the zones were measured. The growth medium and incubation temperature of the indicator bacteria are listed in Table I. Stock cultures in appropriate medium with 20% glycerol were stored at -20°C.

The protein nature of antimicrobial substances produced by isolates was determined by proteinase K enzyme assay. The 20 µl proteinase K (20 mg/ml) was dropped near 1 cm of the colonies and then the plates were incubated for 1 h at 37°C. Finally, indicator bacteria were poured onto plate with 5 ml soft agar and incubated at an optimum growth temperature for the indicators. After the incubation period, diminishing of inhibition zones shaped like a half-moon was considered as an indication that the antimicrobial agent produced by *Enterococcus* sp. isolates was affected by the enzyme proteinase K and antimicrobial substances were recognized as bacteriocin (enterocin).

**Genomic DNA extraction.** Genomic DNA's of bacteriocin producing isolates were extracted from 0.5 ml of overnight cultures using the method of Cancilla *et al.* (1992). DNA precipitates were resuspended in 50 µl Tris-EDTA buffer (pH 8.0).

**Identification of enterocin producer isolates.** 16S rDNA gene sequence of the isolates were amplified with following specific primers: pA (forward) 5'-AGA GTT TGA TCC TGG CTC AG-3' and pE' (reverse) 5'-CCG TCA ATT CCT TTG AGT TT-3' (Edwards *et al.*, 1989) using in a programmable DNA thermocycler (Techne TC3000, Cambridge, U.K.). Fifty microliter PCR reaction mixture consisted of 3 µl of the bacterial DNA solution, 1 µl of each primer, 20 µl nuclease-free water and 25 µl PCR master mix (Fermentas, Vilnius, Lithuania) and following PCR conditions were ensured: initial denaturation cycle at 94°C for 2 min, next 30 cycles of denaturation 94°C for 30 s, annealing at 55°C for 60 s and elongation at 72°C for 90 s, and a final extension cycle at 72°C for 10 min. Amplification products were separated by electrophoresis on 1% agarose (w/v) gels at 85 V for 1.5 h in Tris-acetate-EDTA buffer. Agarose gels were stained with ethidium bromide (20 µg/ml) and photographed under ultraviolet light using a Nikon D5100 digital camera (Nikon Corp., Japan). The size of amplicons was determined by comparison with O'GeneRuler 100-bp DNA ladder (Fermentas). Sequencing of the 16S rDNA gene was performed in RefGen (ODTU Technocity, Ankara, Turkey). The 16S rDNA homology searches were carried out using BLAST software against the NCBI (National Center for Biotechnology Information, Bethesda, MD, USA) database.

Table I  
Growth medium and incubation temperatures of indicator bacteria, and inhibitory spectrum of *Enterococcus* sp. strains

Indicator strains	Growth medium* and incubation temperature	Inhibition zone (Ø mm)										
		MBE 1-9	MBE 12-3	MBE 15-2	MBE 16-5	MBE 20-5	MBE 22-2	MBE 27-2	MBE 27-3	MBE 29-1	MBE 29-3	MBE 31-4
<i>Lactobacillus plantarum</i> LMG2003	MRS, 37°C	-	-	-	-	-	-	-	-	-	-	-
<i>Lactococcus lactis</i> subsp. <i>lactis</i> 1	GM17, 30°C	-	-	-	-	-	-	-	-	-	18	-
<i>Lactococcus lactis</i> subsp. <i>lactis</i> 731	GM17, 30°C	7	9	12	16	-	-	13	10	-	10	-
<i>Lactococcus lactis</i> subsp. <i>lactis</i> T1	GM17, 30°C	16	13	14	-	-	-	14	13	-	15	-
<i>Lactococcus lactis</i> subsp. <i>lactis</i> 105	GM17, 30°C	-	-	-	-	-	-	-	-	-	-	-
<i>Listeria innocua</i> LMG2813	LB, 30°C	7	9	8	6	4	5	9	9	4	6	4
<i>Listeria monocytogenes</i> ATCC15813	LB, 30°C	4	13	17	7	3	5	13	13	3	7	4
<i>Listeria monocytogenes</i> ATCC19115	LB, 30°C	13	5	10	9	6	7	10	6	4	14	6
<i>Listeria monocytogenes</i> ATCC7644	LB, 30°C	12	23	24	10	7	9	26	23	6	11	6
<i>Staphylococcus aureus</i> ATCC25923	TSB, 37°C	7	7	10	9	6	7	9	7	6	8	8
<i>Staphylococcus aureus</i> FRI1003022	TSB, 37°C	7	6	6	5	4	6	7	6	5	6	4
<i>Staphylococcus carnosus</i> LMG2709	TSB, 37°C	8	5	5	6	5	6	7	6	5	15	6
<i>Staphylococcus aureus</i> ATCC29213	TSB, 37°C	11	13	10	15	12	11	10	12	10	12	14
<i>Enterococcus faecalis</i> ATCC29212	GM17, 37°C	8	14	10	6	-	5	8	14	5	8	6
<i>Enterococcus faecalis</i> LMG2708	GM17, 37°C	-	10	-	-	-	-	-	8	-	-	-
<i>Enterococcus faecalis</i> LMG2602	GM17, 37°C	-	15	-	-	-	-	-	-	-	13	-
<i>Bacillus cereus</i> ATCC10876	TSB, 37°C	6	7	5	7	6	6	6	7	5	6	6
<i>Bacillus cereus</i> LMG2732	TSB, 37°C	17	5	16	-	-	-	16	-	-	18	-
<i>Salmonella</i> Typhimurium SL1344	LB, 37°C	10	15	16	14	14	9	12	13	14	17	8
<i>Salmonella</i> Enteritidis ATCC13076	LB, 37°C	7	7	8	7	8	5	16	6	5	7	5
<i>Salmonella</i> Typhimurium ATCC14028	LB, 37°C	-	8	4	4	4	6	-	8	-	-	5
<i>Escherichia coli</i> LMG3083 CFAI (ETEC)	LB, 37°C	5	6	6	4	-	-	6	7	-	7	-
<i>Pediococcus pentosaceus</i> LMG2001	TSB, 37°C	19	15	16	-	-	-	17	14	5	18	-

\* MRS: de Man Rogosa Sharpe, GM17: M17 with glucose (5%), LB: Luria Bertani, TSB: Tryptic Soy Broth

Table II  
PCR primers, annealing temperatures and product sizes for detection of bacteriocin\*, virulence\*\* and amino acid decarboxylase\*\*\* genes.

Gene	Primer sequence (5' to 3')	Annealing temperature (°C)	Product size (bp)	Reference
*entA	AAATATTATGGAAATGGAGTGTAT GCACTTCCCTGGAATTGCTC	56	126	Yousif <i>et al.</i> 2005
*entB	GAAAATGATCACAGAATGCCT A GTTGCATTTAGAGTATACATTTG	50	162	Yousif <i>et al.</i> 2005
*entP	TATGGTAATGGTGT'TTAT'TGTAAT ATGTCCCATACTGCCAAAC	50	120	Yousif <i>et al.</i> 2005
*entL50A/B	TGGGAGCAATCGCAAAATTAG ATTGCCCATCCTTCTCCAAT	52	98	Ben Belgacem <i>et al.</i> 2010
*bac31	TATTACGGAAATGGTTTATAT'TGT TCTAGGAGCCCAAGGGCC	50	123	Yousif <i>et al.</i> 2005
*entAS48	GAGGAGT'TTCATGAT'TTAAAGA CATATTGTTAAAT'TACCAAGCAA	50	340	Yousif <i>et al.</i> 2005
*entQ	ATGAATTTTCTTCTTAAAAATGGTATCGCA TTAACAAGAAATTTT'TCCCATGGCAA	56	105	Ben Belgacem <i>et al.</i> 2010
*ent1071A/B	CCTATTGGGGGAGAGTCGGT ATACATTCTTCCACTTATTTTT	51	343	Ben Belgacem <i>et al.</i> 2010

Table II continued

Gene	Primer sequence (5' to 3')	Annealing temperature (°C)	Product size (bp)	Reference
* <i>cyl<sub>L/S</sub></i>	GTGTTGAGGAAATGGAAGCG TCTCAGCCTGAACATCTCCAC	60	324	Brandao <i>et al.</i> 2010
* <i>entX</i>	GTTTCTGTAAAAGAGATGAAAC CCTCTTAATCATTAACCATAC	50	500	Edalatian <i>et al.</i> 2012
** <i>efaA<sub>fm</sub></i>	AACAGATCCGCATGAATA CATTTTCATCATCTGATAGTA	54	735	Reviriego <i>et al.</i> 2005
** <i>efaA<sub>fs</sub></i>	GACAGACCCTCACGAATA AGTTCATCATGCTGTAGTA	54	705	Reviriego <i>et al.</i> 2005
** <i>ccf</i>	GGGAATTGAGTAGTGAAGAAG AGCCGCTAAAATCGGTAAAAT	54	543	Reviriego <i>et al.</i> 2005
** <i>cpd</i>	TGGTGGGTTATTTTTCAATTC TACGGCTCTGGCTTACTA	54	782	Reviriego <i>et al.</i> 2005
** <i>cob</i>	AACATTCAGCAAACAAAGC TTGTCATAAAGAGTGGTCA	54	1405	Reviriego <i>et al.</i> 2005
** <i>esp<sub>fm</sub></i>	TTGCTAATGCAAGTCACGTCC GCATCAACACTTGCATTACCGAA	54	955	Reviriego <i>et al.</i> 2005
** <i>esp<sub>fs</sub></i>	TTGCTAATGCTAGTCCACGACC GCGTCAACACTTGCATTGCCGAA	54	933	Reviriego <i>et al.</i> 2005
** <i>ace</i>	AAAGTAGAATTAGATCCACAC TCTATCACATTTCGGTTGCG	54	350	Ben Belgacem <i>et al.</i> 2010
** <i>gelE</i>	ACCCCGTATCATTGGTT ACGCATTGCTTTTCCATC	54	419	Reviriego <i>et al.</i> 2005
** <i>cad</i>	TGCTTTGTCATTGACAATCCG ACTTTTTCCCAACCCCTCAA	54	1299	Reviriego <i>et al.</i> 2005
** <i>agg</i>	AAGAAAAAGAAGTAGACCAAC AAACGGCAAGACAAGTAAATA	56	1553	Eaton and Gasson 2001
** <i>cylA</i>	TGGATGATAGTGATAGGAAGT TCTACAGTAAATCTTTCGTCA	54	517	Reviriego <i>et al.</i> 2005
** <i>cylB</i>	ATTCCTACCTATGTTCTGTTA AATAAACTCTTCTTTCCAAC	54	843	Reviriego <i>et al.</i> 2005
** <i>cylM</i>	CTGATGGAAAGAAGATAGTAT TGAGTTGGTCTGATTACATTT	54	742	Reviriego <i>et al.</i> 2005
*** <i>hdc</i>	GGNATNGTNWSNTAYGAYMGNGCNGA ATNGCDATNGCNSWCCANACNCCRTA	53	372	de las Rivas <i>et al.</i> 2006
*** <i>tdc</i>	TGGYTNGTNCNCARACNAARCACTA ACRTARTCNACCATRTTRAARTCNGG	53	825	de las Rivas <i>et al.</i> 2006
*** <i>odc</i>	TWYMAYGCNGAYAARACNTAYTTYGT ACRCANAGNACNCCNGGNGGRTANGG	53	1440	de las Rivas <i>et al.</i> 2006
*** <i>ldc</i>	CAYRTNCCNNGNCAYAA GGDATNCCNNGNGGRTA	53	1185	de las Rivas <i>et al.</i> 2006

PCR primers, annealing temperatures and product sizes for detection of bacteriocin\*, virulence\*\* and amino acid decarboxylase\*\*\* genes  
Y: C or T; R: A or G; W: A or T; S: C or G; M: A or C; D: A, G or T.

**Detection of enterocin genes.** The presence of enterocin-encoding genes was detected according to Edalatian *et al.* (2012), by using the most common enterocin primers. Interaction primers, annealing temperatures and the size of amplicons are given in Table II. PCR for enterocin genes were performed using the following parameters: initial denaturation at 94°C for 5 min, 35 cycles of denaturation at 94°C for 1 min, annealing at an appropriate temperature for 1 min, and elongation at 72°C for 40 s and final exten-

sion at 72°C for 10 min. PCR products were analyzed by agarose gel (1.5%, w/v) electrophoresis with reference to O'GeneRuler 100-bp DNA ladder (Fermentas), the bands stained with ethidium bromide and photographed with a Nikon D5100 digital camera under ultraviolet light. *E. faecium* EYT31 strain (*entA*<sup>+</sup>, *entB*<sup>+</sup> and *entP*<sup>+</sup>) obtained from Özden-Tuncer *et al.* (2013), was used as positive control.

**Cross-protection activity test.** The purpose of this assay was to detect the susceptibility of each strain to

antimicrobial substances produced by all others. Consequently, enterococci strains were used as producers and indicators. Producer bacteria were spotted on appropriate solid media and then enterocin producer strains were poured onto plates with soft MRS agar as indicators. After incubation at 37°C for 18 h inhibition zones were examined.

**Haemolytic and gelatinase activity.** *Enterococcus* sp. strains were grown in MRS broth medium at 37°C for 18 h and then streaked onto Columbia Agar (Laboratorios Conda S.A., Madrid, Spain) containing 5% defibrinated sheep blood for determining haemolytic activity. After an incubation period (37°C for 48 h) presence of zones of clearing around the colonies were interpreted as  $\beta$ -haemolysis. The absence of zones around the colonies or green zones around the colonies was interpreted as  $\gamma$ -haemolysis and  $\alpha$ -haemolysis, respectively (Cariolato *et al.*, 2008). Gelatinase production abilities of the enterocin producer strains were investigated on Todd-Hewitt agar presence of gelatinase (30 g/l). Briefly, strains were grown in MRS broth medium (at 37°C for 24 h) then streak onto Todd-Hewitt agar (Liofilchem, Roseto degli Abruzzi, Italy). After incubation at 37°C for 24 h, petri dishes were placed 4°C for 5 h before examination for the zone of turbidity around the colonies, indicating hydrolysis (Eaton and Gasson, 2001).

**Antibiotic susceptibility.** Antibiotic susceptibility of enterocin producer *E. faecium* and *E. faecalis* strains were determined by disc diffusion method on Mueller Hinton agar (Lab M) plates according to the recommendation of Clinical and Laboratory Standards Institute (CLSI, 2012). The antibiotics used in this study were as follows: doxycycline (30  $\mu$ g), norfloxacin (10  $\mu$ g), vancomycin (30  $\mu$ g), chloramphenicol (30  $\mu$ g), rifampicin (5  $\mu$ g), minocycline (30  $\mu$ g), penicillin (10  $\mu$ g), streptomycin (300  $\mu$ g), levofloxacin (5  $\mu$ g), ciprofloxacin (5  $\mu$ g), ampicillin (10  $\mu$ g), quinupristin-dalfopristin (15  $\mu$ g), nitrofurantoin (300  $\mu$ g), tetracycline (30  $\mu$ g), gentamicin (120  $\mu$ g), erythromycin (15  $\mu$ g), linezolid (30  $\mu$ g) and teicoplanin (30  $\mu$ g).

**Presence of *vanA* and *vanB* genes.** *E. faecium* and *E. faecalis* strains were searched for the presence of *vanA* and *vanB* genes with the following primers: VanA1 [5'-GGG AAA ACG ACA ATT GC-3'] and VanA2 [5'-GTACAA TGC GGC CGT TA-3'] for *vanA* gene (Dutka-Malen *et al.*, 1995) and VanB [5'-GTG CTG CGA GAT ACC ACA GA-3'] and VanBrev [5'-CGA ACA CCA TGC AAC ATT TC-3'] for *vanB* gene (Reviriego *et al.*, 2005). Product size of 732 bp and 1,145 bp were screened for *vanA* and *vanB* genes, respectively. PCR conditions were performed using the following cycling parameters: initial denaturation cycle at 94°C for 2 min, next 30 cycles of denaturation 94°C for 1 min, annealing at 54°C for 1 min and elongation at

72°C for 1 min, and a final extension cycle at 72°C for 10 min. Electrophoresis of the amplification products was performed on 1.5% (w/v) agarose gel.

**Detection of virulence factors.** Presence of the *gelE* (gelatinase), *efaA<sub>fm</sub>* and *efaA<sub>fs</sub>* (cell wall adhesin for *E. faecium* and *E. faecalis*, respectively), *cpd*, *cob*, *ccf* and *cad* (sex pheromones), *ace* (collagen adhesin), *esp<sub>fm</sub>* and *esp<sub>fs</sub>* (cell wall-associated protein in *E. faecium* and *E. faecalis*, respectively), *agg* (aggregation substance), *cylM*, *cylB* and *cylA* (cytolysin) genes were investigated by PCR. PCR primers, annealing temperatures and product sizes for detection of virulence factors are listed in Table II. PCR conditions were performed using the following cycling parameters: initial denaturation cycle at 95°C for 5 min, next 35 cycles of denaturation 95°C for 30 s, annealing at an appropriate temperature for 30 s and elongation at 72°C for 1 min, and a final extension cycle at 72°C for 10 min. Electrophoresis of the amplicons was performed on 1.5% (w/v) agarose gel.

**Detection of decarboxylase activity and amino acid decarboxylase genes.** The amino acid decarboxylase activity of *Enterococcus* sp. strains was estimated according to the method proposed by Bover-Cid *et al.* (1999). Briefly 5 g of amino acid precursors (histidine, lysine, ornithine and tyrosine (Merck) was added to 1 l decarboxylase screening medium. Enterococcal cultures were spotted onto these plates with and without amino acids and incubated at 37°C for 2–5 days. After incubation time, the decarboxylation activity of the *Enterococcus* sp. strains was determined phenotypically by yellow to purple color changes on the petri dishes. Tyraminogenic *E. faecium* NYE54 strain obtained from Inoglu and Tuncer (2013) was used as the positive control.

PCR amplification of histidine, lysine, ornithin and tyrosine decarboxylase genes (*hdc*, *ldc*, *odc* and *tdc*, respectively) were carried out based on the method of de las Rivas *et al.* (2012). PCR primers, annealing temperatures and product sizes for detection of amino acid decarboxylase genes are listed in Table II. PCR conditions were performed using the following cycling parameters: initial denaturation cycle at 95°C for 10 min, next 30 cycles of denaturation 95°C for 30 s, annealing at an appropriate temperature for 30 seconds and elongation at 72°C for 2 min, and a final extension cycle at 72°C for 20 min. Electrophoresis of the amplicons was performed on 1.5% (w/v) agarose gel. *E. faecium* NYE54 strain (*tdc*<sup>+</sup>) (Inoglu and Tuncer, 2013) was used as the positive control.

## Results

**Isolation and phenotypic identification of *Enterococcus* sp. strains.** A total of 100 presumptive *Enterococcus* strains isolated from 33 different cheese samples



Table IV  
Cross protection activity of the enterocin producer *Enterococcus* sp. strains.

Producing strains	Indicator strains										
	MBE 1-9	MBE 12-3	MBE 15-2	MBE 16-5	MBE 20-5	MBE 22-2	MBE 27-2	MBE 27-3	MBE 29-1	MBE 29-3	MBE 31-4
<i>E. faecalis</i> MBE1-9	-	+	+	+	-	+	+	+	+	-	+
<i>E. faecium</i> MBE12-3	-	-	-	+	+	+	-	-	+	+	+
<i>E. faecium</i> MBE15-2	-	-	-	+	-	+	-	-	+	-	+
<i>E. faecium</i> MBE16-5	-	-	-	-	-	-	-	-	+	-	+
<i>E. faecalis</i> MBE20-5	-	-	-	-	-	-	-	-	-	-	-
<i>E. faecium</i> MBE22-2	-	-	-	-	+	-	-	-	+	-	-
<i>E. faecium</i> MBE27-2	-	-	-	+	+	+	-	-	+	-	+
<i>E. faecium</i> MBE27-3	-	-	-	+	+	+	-	-	+	-	+
<i>E. faecalis</i> MBE29-1	-	-	-	-	-	-	-	-	-	-	-
<i>E. faecalis</i> MBE29-3	-	+	+	+	+	+	-	-	+	-	+
<i>E. faecalis</i> MBE31-4	-	-	-	-	-	-	-	-	-	-	-

Table V  
Antibiotic susceptibility of enterocin producer *Enterococcus* sp. strains.

Strains	Antibiotics*																	
	S	CN	MH	TEC	DO	CIP	NOR	QD	RD	VA	TE	P	E	AMP	F	LZD	C	LEV
<i>E. faecalis</i> MBE1-9	R <sup>2</sup>	S	S	S	S	S	I	R	S	S	I	S	I	S	S	S	R	S
<i>E. faecium</i> MBE12-3	S	S	S	S	S	I	S	S	R	S	S	S	I	S	S	S	S	S
<i>E. faecium</i> MBE15-2	S	S	S	S	S	R	R	S	R	S	S	R	I	S	S	S	S	S
<i>E. faecium</i> MBE16-5	S	S	S	S	S	I	S	I	R	S	S	R	R	S	S	S	S	S
<i>E. faecalis</i> MBE20-5	S	S	I	S	S	S	S	S	R	S	R	S	I	S	S	S	S	S
<i>E. faecium</i> MBE22-2	S	S	S	S	S	I	S	I	R	S	S	S	R	S	S	S	S	S
<i>E. faecium</i> MBE27-2	S	S	S	S	S	S	S	S	R	S	S	S	I	S	S	S	S	S
<i>E. faecium</i> MBE27-3	S	S	S	S	S	R	I	S	R	S	S	S	I	S	S	S	S	S
<i>E. faecalis</i> MBE29-1	S	S	S	S	S	S	S	R	I	I	S	S	S	S	S	S	S	S
<i>E. faecalis</i> MBE29-3	S	S	S	S	S	S	S	R	I	S	S	S	I	S	S	S	S	S
<i>E. faecalis</i> MBE31-4	S	S	S	S	S	I	S	R	R	S	S	S	S	S	S	S	S	S

\* DO: doxycycline (30 µg), NOR: norfloxacin (10 µg), VA: vancomycin (30 µg), C: chloramphenicol (30 µg), RD: rifampicin (5 µg), MH: minocycline (30 µg), P: penicillin (10 µg), S: streptomycin (300 µg), LEV: levofloxacin (5 µg), CIP: ciprofloxacin (5 µg), AMP: ampicillin (10 µg), QD: quinupristin-dalfopristin (15 µg), F: Nitrofurantoin (300 µg), TE: tetracycline (30 µg), CN: gentamicin (120 µg), E: erythromycin (15 µg), LZD: linezolid (30 µg), TEC: teicoplanin (30 µg). Susceptibilities of *Enterococcus* sp. strains were determined according to CLSI 2012. <sup>2</sup>R: Resistant, I: Intermediary, S: sensitive.

**Presence of *vanA* and *vanB*.** *E. faecalis* MBE1-9, *E. faecalis* MBE20-5, *E. faecalis* MBE29-1 and *E. faecalis* MBE29-3 strains carry the *vanA* gene among the 11 enterocin producer. Remaining enterocin producers do not contain both *vanA* and *vanB* genes (Table VI).

**Detection of virulence factors.** The *efaA<sub>fs</sub>*, *cad* and *cyl<sub>M</sub>* genes were not detected in any strains while *ccf* and *cylB* genes were amplified by all the strains. The *esp<sub>fm</sub>* gene was found only in *E. faecalis* MBE20-5. *esp<sub>fs</sub>* was found in *E. faecalis* MBE20-5 and MBE31-4 and *agg* was found in *E. faecalis* MBE1-9 and MBE29-3, respectively. *gelE* gene was detected in all *Enterococcus* strains except *E. faecium* MBE22-2. *cpd*, *cob*,

*ace* and *cylA* were also found in several isolates. Virulence traits of all *Enterococcus* sp. strains are given in Table VI.

**Detection of decarboxylase activity and amino acid decarboxylase genes.** The isolated DNA of the *Enterococcus* sp. strains was subjected to PCR amplification to detect the presence of the histidine, lysine, ornithin and tyrosine decarboxylase genes (*hdc*, *ldc*, *odc* and *tdc*, respectively). All of the enterocin producer strains except *E. faecalis* MBE1-9, have phenotypically tyrosine decarboxylase activity and *tdc* gene was also detected in the same strains. The *hdc*, *ldc* and *odc* genes were not detected in any strains (Table VI).

Table VI  
Presence of virulens factors, vancomycin and biogenic amine genes in *Enterococcus* sp. strains.

Strains	Virulens factors*													Vancomycin**			Biogenic amine***			
	<i>efaA<sub>fm</sub></i>	<i>efaA<sub>fs</sub></i>	<i>ccf</i>	<i>cpd</i>	<i>cob</i>	<i>cad</i>	<i>esp<sub>fm</sub></i>	<i>esp<sub>fs</sub></i>	<i>ace</i>	<i>gelE</i>	<i>agg</i>	<i>cylA</i>	<i>cylB</i>	<i>cylM</i>	<i>vanA</i>	<i>vanB</i>	<i>hdc</i>	<i>tdc</i>	<i>odc</i>	<i>ldc</i>
<i>E. faecalis</i> MBE1-9	-	-	+	+	+	-	-	-	+	+	+	+	+	-	+	-	-	-	-	-
<i>E. faecium</i> MBE12-3	+	-	+	-	-	-	-	-	-	+	-	-	+	-	-	-	-	+	-	-
<i>E. faecium</i> MBE15-2	+	-	+	-	-	-	-	-	-	+	-	-	+	-	-	-	-	+	-	-
<i>E. faecium</i> MBE16-5	-	-	+	+	-	-	-	-	-	+	-	-	+	-	-	-	-	+	-	-
<i>E. faecalis</i> MBE20-5	+	-	+	+	+	-	+	+	+	+	-	+	+	-	+	-	-	+	-	-
<i>E. faecium</i> MBE22-2	+	-	+	-	-	-	-	-	-	-	-	-	+	-	-	-	-	+	-	-
<i>E. faecium</i> MBE27-2	+	-	+	+	-	-	-	-	-	+	-	+	+	-	-	-	-	+	-	-
<i>E. faecium</i> MBE27-3	+	-	+	+	-	-	-	-	-	+	-	-	+	-	-	-	-	+	-	-
<i>E. faecalis</i> MBE29-1	+	-	+	+	+	-	-	-	+	+	-	+	+	-	+	-	-	+	-	-
<i>E. faecalis</i> MBE29-3	-	-	+	+	+	-	-	-	+	+	+	+	+	-	+	-	-	+	-	-
<i>E. faecalis</i> MBE31-4	-	-	+	+	-	-	-	+	+	+	-	-	+	-	-	-	-	+	-	-

\**efaA<sub>fm</sub>* and *efaA<sub>fs</sub>*: cell wall adhesin (from *E. faecium* and *E. faecalis*, respectively); *ccf*, *cpd*, *cob* and *cad*: sex pheromone; *esp<sub>fm</sub>* and *esp<sub>fs</sub>*: enterococcal surface protein (from *E. faecium* and *E. faecalis*, respectively); *ace*: collagen adhesion; *gelE*: gelatinase; *agg*: aggregation substance; *cylM*, *cylB*, and *cylA*: cytolyisin.

\*\**vanA*: vancomycin A structural gene; *vanB*: vancomycin B structural gene.

\*\*\**hdc*: histidine decarboxylase gene; *tdc*: tyrosine carboxylase gene; *odc*: ornithin decarboxylase gene; *ldc*: lysin decarboxylase gene.

## Discussion

Enterococci easily adapt to several food systems due to their high salt and acid tolerance. It is believed that these bacteria contribute to the flavor and the aroma formation of some different European and Mediterranean cheeses. *E. faecium*, *E. faecalis* and *E. durans* species were frequently isolated from traditional dairy products (Giraffa, 2002; 2003; Martin-Platero *et al.*, 2009; Yogurtcu and Tuncer, 2013; Furlaneto-Maia *et al.*, 2014). In the present study a total of 100 *Enterococcus* strains were isolated from various artisanal cheese samples and 11 strains were determined as enterocin producers and also characterized on species level as *E. faecium* (6 strains) and *E. faecalis* (5 strains). The diversity of *Enterococcus* species from cheese depends on the ecological niche that they were isolated from (Rivas *et al.*, 2012). Similar to other studies we were found that *E. faecium* and *E. faecalis* were mostly isolated enterococcal strains from traditional Turkish cheeses (Tuncer, 2009; Yogurtcu and Tuncer, 2013).

Some *Enterococcus* sp. strains show inhibition activity with their enterocins against food spoilage and food pathogen microorganisms (Khan *et al.*, 2010; Javed *et al.*, 2011; Özden-Tuncer *et al.*, 2013). It was observed that all *E. faecium* and *E. faecalis* strains showed inhibition zones to all *Listeria* sp. and *Staphylococcus* sp. strains used in this study besides closely related species. On the other hand, inhibition zones against 2 *Salmonella* and 1 *E. coli* strains were also determined.

Garcia-Cano *et al.* (2014), indicated that 1 *E. faecium* and 1 *E. faecalis* strains isolated from Cotija cheese displayed inhibitor activity against *S. enterica* Typhimurium, *E. coli* besides *L. monocytogenes* and *S. aureus* similar to our findings.

*E. faecalis* MBE29-1 and *E. faecalis* MBE29-3 strains were found to harbor only *entX* structural gene while *E. faecium* MBE16-5 and *E. faecalis* MBE31-4 strains carried *entA* and *entX*; *E. faecium* MBE15-2 and *E. faecium* MBE 27-3 strains carry *entA*, *entB* and *entX*; *E. faecium* MBE12-3 strain carried *entA* and *entB*; *E. faecium* MBE22-2 strain carried *entA*, *entP* and *entX* structural genes. *E. faecium* MBE27-2 was identified as the only producer strain that harbored 4 structural genes (*entA*, *entB*, *entP* and *entX*) at the same time. It is not unusual that multiple structural enterocin genes may be present in one strain and it is indicated that many *Enterococcus* sp. strains produce more than one bacteriocin similar to our results (Edalatian *et al.*, 2012; Özden-Tuncer *et al.*, 2013; Rehaïem *et al.*, 2014). On the other hand, *E. faecalis* MBE1-9 and MBE20-5 strains have not given the PCR product with the most common enterocin primers tested in this study suggesting that bacteriocins produced by these strains could be new enterocins. Further analyses should be needed for characterization of these enterocins.

However, many *Enterococcus* sp. strains adapt to food fermentation processes as a starter or as a protective culture, they may harbor antibiotic resistance and some virulence determinants that increase the



risk of pathogenicity of these strains (Martin-Platero *et al.*, 2009; Edalatian *et al.*, 2012). In the present study, 11 enterocin producer strains were found to be resistant to one or more antibiotics. All strains showed sensitivity against vancomycin but four of them were carrying *vanA* genes. Furlaneto-Maia *et al.* (2014), reported that 5 *Enterococcus* sp. isolates harbored *vanA* gene, while they were susceptible to vancomycin phenotypically. In addition the same results for ampicillin, tetracycline and linezolid susceptibilities were attained in the study of Pesavento *et al.* 2014. We determined that all enterocin producer strains showed susceptibility against gentamicin, teicoplanin, doxycycline, ampicillin, nitrofurantoin, linezolid and levofloxacin. On the other hand, most of the resistance in the strains obtained against rifampicin and quinupristin-dalfopristin similar to other results obtained in *Enterococcus* sp. strains isolated from Turkish Tulum cheese (Özden-Tuncer *et al.*, 2013).

Only *E. faecalis* MBE29-3 showed gelatinase activity on Todd-Hewitt agar but all the enterocin producers strains except *E. faecium* MBE 22-2 had gene *gelE*. These results suggest that in strains which have no gelatinase activity phenotypically associated with *gelE* the gene was silent or expressed at a low level (Lopes *et al.*, 2006). Similar observations were indicated for clinical and food originated *Enterococcus* isolates (Cariolato *et al.*, 2008; Ben Belgacem *et al.*, 2010; Özden-Tuncer *et al.*, 2013; Inoğlu and Tuncer, 2013; Tuncer *et al.*, 2014; Medeiros *et al.*, 2014). In addition, many of the virulence genes were detected in enterocin producer strains. PCR analysis revealed that all strains contained at least 3 virulence genes. However, sex pheromone genes (*cpd*, *cob*, *ccf* and *cad*) were not accepted as virulence factors their production in *Enterococcus* sp. strains are assume that dissamination of virulence factors, and these genes were detected frequently in *Enterococcus* sp. isolates from cheese (Eaton and Gasson, 2001; Valenzuela *et al.*, 2008; Ben Belgacem *et al.*, 2010; Inoğlu and Tuncer, 2013). Martin *et al.* (2006), confirmed that *efaA<sub>fs</sub>* and *efaA<sub>fm</sub>* were only found in *E. faecalis* and *E. faecium*, respectively. However, Barbosa *et al.* (2010) and Inoğlu and Tuncer (2013) were indicated that *efaA<sub>fs</sub>* gene found in *E. faecium* and *efaA<sub>fm</sub>* was found in *E. faecalis* strain, as confirmed in this study. The *agg* gene product mediates the establishment of contact between bacteria and host cells and facilitating colonization, therefore *agg* gene harboring strains were unwanted in food. The *agg* gene was detected only in *E. faecalis* strains by other researchers (Eaton and Gasson, 2001; Franz *et al.*, 2001; Martin *et al.*, 2006; Inoğlu and Tuncer, 2013), as confirmed in this study. However, Barbosa *et al.* (2010), showed that *agg* gene may be found in *E. faecium* strains too. The *ace* gene was also having an important role in colonization

like *agg* gene and it is reported that *Enterococcus* sp. isolates containing *ace* gene were found to be positively correlated with biofilm formation (Singh *et al.*, 2010, Medeiros *et al.*, 2014). In our study, the strains containing the *agg* and *ace* genes individually and the strains possessing also both genes should be investigated for biofilm formation in a further study.

Cytolysin is a bacterial toxin involved in the formation of haemolytic activity. Several studies confirmed that *Enterococcus* sp. strains harboring cytolysin genes may have  $\beta$ -haemolytic activity (Eaton and Gasson, 2001; Theppangna *et al.*, 2007). However, different researchers found that, non  $\beta$ -haemolytic strains may carry the cytolysin genes similar to the present study (Cosentino *et al.*, 2010; Medeiros *et al.*, 2014). Medeiros *et al.* (2014), showed that *Enterococcus* sp. isolate originated from food had non  $\beta$ -hemolytic activity, but carried *cylA* gene, as confirmed in this study. It is also reported that in non haemolytic strains cytolysin determinant may behave as silent. On the other hand, the lack of a correlation between phenotypic and genotypic cytolysin production may suggest some genetic rearrangements in *cyl* operon like missing genes (Semedo *et al.*, 2003, Gaspar *et al.*, 2009). In our study *cylB* gene was detected in all strains while *cylA* gene was detected only in 5 strains. However, *cylM* was not detected in any strains and in all strains the lack of  $\beta$ -haemolytic activity was determined.

Another undesirable case for *Enterococcus* sp. strains is biogenic amine production ability (Valenzuela *et al.*, 2008). In the present study, tyrosine decarboxylating ability was determined in 10 *Enterococcus* sp. strains phenotypically and gene *tdc* was also detected in the same strains. Only *E. faecalis* MBE1-9 has no decarboxylating tyrosine does not harbor *tcd* gene as well. Similar to our results, several researchers indicated that tyramine was frequently produced by *Enterococcus* sp. strains (Tuncer, 2009; Komprda *et al.*, 2010; Kalhotka *et al.*, 2012; Inoğlu and Tuncer, 2013). PCR study showed compatibility with the result of the phenotypically tyrosine decarboxylating ability, as confirmed by Inoğlu and Tuncer (2013) and Yüceer and Özden-Tuncer (2015).

## Conclusion

Many of the enterococci originating from food may produce enterocins are active against pathogenic or spoilage microorganisms, particularly *Listeria* spp. and *Staphylococcus aureus*. In the present study enterocins were identified using known enterocin primers and multiple enterocin genes were detected in 7 strains. Two enterocins could not be identified and need to be investigated whether they are new bacteriocins.

However, all enterocin producers were found to have several virulence factors such *i.e.* *gelE*, *agg*, *cpd*, *cob*, *ace*, *cylA* and *tdc* genes. In addition 6 strains had multiple antibiotic resistance patterns. Lack of hemolytic and gelatinase activity (except *E. faecalis* MBE29-3) in bacteriocin producing strains are advantageous, although their virulence characteristics make them risky in terms of consumer health.

#### Acknowledgements

This research was financially supported by the Süleyman Demirel University, Scientific Research Project Coordination Unit through Project No. 3819-YL2-13.

#### Literature

- Barbosa J., P.A. Gibbs and P. Teixeira.** 2010. Virulence factors among enterococci isolated from traditional fermented meat products produced in the North of Portugal. *Food Control* 21: 651–656.
- Ben Belgacem Z., H. Abriouel, N. Ben Omar, R. Lucas, M. Martínez-Canamero, A. Gálvez and M. Manai.** 2010. Antimicrobial activity, safety aspects, and some technological properties of bacteriocinogenic *Enterococcus faecium* from artisanal Tunisian fermented meat. *Food Control* 21: 462–470.
- Bover-Cid S., M. Hugas, M. Izquierdo-Pulido and M.C. Vidal-Carou.** 1999. Amino acid decarboxylase activity of bacteria isolated from fermented pork sausages. *Int. J. Food Microbiol.* 66: 185–189.
- Brandao A. T., Almeida, E. Munoz-Atieza, C. Torres, G. Igrejas, P.E. Hernández, L.M. Cintas, P. Poeta and C. Herranz.** 2010. Antimicrobial activity and occurrence of bacteriocin structural genes in *Snterococcus* spp. of human and animal origin isolated in Portugal. *Arch. Microbiol.* 192: 927–936.
- Cancilla M.R., I.B. Powell, A.J. Hillier and B.E. Davidson.** 1992. Rapid genomic fingerprinting of *Lactococcus lactis* strains by arbitrarily primed polymerase chain reaction with 32P and fluorescent labels. *Appl. Environ. Microbiol.* 58: 1772–1775.
- Cariolato D., C.Andrighetto and A. Lombardi** 2008. Occurrence of virulence factors and antibiotic resistances in *Enterococcus faecalis* and *Enterococcus faecium* collected from dairy and human samples in North Italy. *Food Control* 19: 886–892.
- Chuang O.N., P.M. Schlievert, C.L. Wells, D.A. Manias and T.J. Tripp.** 2009. Multiple functional domains of *Enterococcus faecalis* aggregation substance Asc10 contribute to endocarditis virulence. *Infect. Immun.* 77: 539–548.
- CLSI.** 2012. Clinical and Laboratory Standards Institute, Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard-Eleventh Edition. M02-A11, Vol 32 No. 1.
- Cosentino S., G.S. Podda, A. Corda, M.E. Fadda, M. Deplano and M.B. Pisano.** 2010. Molecular detection of virulence factors and antibiotic resistance pattern in clinical *Enterococcus faecalis* strains in Sardinia. *J. Prev. Med. Hygiene* 51: 31–36.
- de las Rivas B., A. Marcobal, A.V. Carrascosa and R. Munoz.** 2006. PCR detection of foodborne bacteria producing the biogenic amines histamine, tyramine, putrescine and cadaverine. *J. Food Protect.* 69: 2509–2514.
- Dutka-Malen S., S. Evers and P. Courvalin.** 1995. Detection of glycopeptide resistance genotypes and identification to the species level of clinically relevant enterococci by PCR. *J. Clin. Microbiol.* 33: 24–27.
- Eaton T.J. and M.J. Gasson.** 2001. Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates. *Appl. Environ. Microbiol.* 67: 1628–1635.
- Edalatian M.R., M.B.H. Najafi, S.A. Mortazavi, Á. Alegría, S. Delgado, M.R. Bassami and B. Mayo.** 2012. Production of bacteriocins by *Enterococcus* spp. isolated from traditional, Iranian, raw milk cheeses, and detection of their encoding genes. *Eur. Food Res. Technol.* 234: 789–796.
- Edwards U., T. Rogall, H. Blocker, M. Emde and E.C. Bottger.** 1989. Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Research* 17: 7843–7853.
- Franz C.M.A.P., A.B. Muscholl-Silberhorn, N.M.K. Yousif, M. Vancanneyt, J. Swings and W.H. Holzapfel.** 2001. Incidence of virulence factors and antibiotic resistance among enterococci isolated from food. *Appl. Environ. Microbiol.* 67: 4385–4389.
- Franz C.M.A.P., M.E. Stiles, K.H. Schleifer and W.H. Holzapfel.** 2003. Enterococci in foods- a conundrum for food safety. *Int. J. Food Microbiol.* 88: 105–122.
- Furlaneto-Maia L., K.R. Rocha, F.C. Henrique, A.Giazzi and M.C. Furlaneto** 2014. Antimicrobial resistance in *Enterococcus* spp. isolated from soft cheese in southern Brazil. *Adv. Microbiol.* 4: 175–181.
- García-Cano I., C.E. Serrano-Maldonado, M. Olvera-García, E. Delgado-Arciniega, C. Peña-Montes, G. Mendoza-Hernández and M. Quirasco.** 2014. Antibacterial activity produced by *Enterococcus* spp. isolated from an artisanal Mexican dairy product, Cotija cheese. *LWT Food Sci. Technol.* 59: 26–34.
- Gaspar F.B., M.T.B. Crespo, and M.F.S. Lopes.** 2009. Proposal for a reliable enterococcal cytolysin production assay avoiding apparent incongruence between phenotype and genotype. *J. Med. Microbiol.* 58: 1122–1124.
- Giraffa G.** 2002. Enterococci from foods. *FEMS Microbiol. Rev.* 26(2): 163–171.
- Giraffa G.** 2003. Functionality of enterococci in dairy products. *Int. J. Food Microbiol.* 88: 215–222.
- Inoğlu Z.N. and Y. Tuncer.** 2013. Safety assessment of *Enterococcus faecium* and *Enterococcus faecalis* strains isolated from Turkish Tulum Cheese. *J. Food Safety* 33: 369–377.
- Javed A., T. Masud, Q. ul Ain, M. Imran and S. Maqsood.** 2011. Enterocins of *Enterococcus faecium*, emerging natural food preservatives. *Ann. Microbiol.* 61: 699–708.
- Kalhotka L., I. Manga, J. Přichystalová, M. Hůlová, M. Vyletřelová and K. Šustová.** 2012. Decarboxylase activity test of the genus *Enterococcus* isolated from goat milk and cheese. *Acta Veterinaria Brno* 81: 145–151.
- Khan H., S. Flint and P.L. Yu.** 2010. Enterocins in food preservation. *Int. J. Food Microbiol.* 141: 1–10.
- Komprda T., P. Sladková, E. Petirová, V. Dohnal and R. Burdychová.** 2010. Tyrosine- and histidine-decarboxylase positive lactic acid bacteria and enterococci in dry fermented sausages. *Meat Sci.* 86: 870–877.
- Lindenstrauß A.G., M. Pavlovic, A. Bringmann, J. Behr, M.A. Ehrmann and R.F. Vogel.** 2011. Comparison of genotypic and phenotypic cluster analyses of virulence determinants and possible role of CRISPR elements towards their incidence in *Enterococcus faecalis* and *Enterococcus faecium*. *Sys. Appl. Microbiol.* 34: 553–560.
- Lopes M.F.S., A.N. Simões R. Tenreiro, J.J. Marques, M.T.B. Crespo.** 2006. Activity and expression of a virulence factor, gelatinase, in dairy enterococci. *Int. J. Food. Microbiol.* 112: 208–214.
- Manolopoulou E., P. Sarantinopoulos, E. Zoidou, A. Aktypis, E. Moschopoulou, I.G. Kandarakis and E.M. Anifantakis.** 2003. Evolution of microbial populations during traditional Feta cheese manufacture and ripening. *Int. J. Food Microbiol.* 82: 153–161.
- Martín M., J. Gutiérrez, R. Criado, C. Herranz, L.M. Cintas and P.E. Hernández.** 2006. Genes encoding bacteriocins and their expres-

- sion and potential virulence factors of enterococci isolated from wood pigeons (*Columba palumbus*). *J. Food Protect.* 69: 520–531.
- Martin-Platero A.M., E. Valdivia, M. Maqueda and M. Martinez-Bueno.** 2009. Characterization and safety evaluation of enterococci isolated from Spanish goats' milk cheeses. *Int. J. Food Microbiol.* 132: 24–32.
- Medeiros A.W., R.I. Pereira, D.V. Oliveira, P.D. Martins, P.A. d'Azevedo, S. Van der Sand, J. Frazzon, A.P.G. Frazzon.** 2014. Molecular detection of virulence factors among food and clinical *Enterococcus faecalis* strains in south Brazil. *Braz. J. Microbiol.* 45(1): 327–332.
- Moreno M.R.F., P. Sarantinopoulos, E. Tsakalidou and L. De Vuyst.** 2006. The role and application of enterococci in food and health. *Int. J. Food Microbiol.* 106: 1–24.
- Nes I.F., D.B. Diep and H. Holo.** 2007. Bacteriocin diversity in *Streptococcus* and *Enterococcus*. *J. Bacteriol.* 189 (4): 1189–1198.
- Özden-Tuncer B., Z. Ay and Y. Tuncer.** 2013. Occurrence of enterocin genes, virulence factors and antibiotic resistance in three bacteriocin producer *Enterococcus faecium* strains isolated from Turkish tulum cheese. *Turkish J. Biol.* 37: 443–449.
- Pesavento G., C. Calonico, B. Ducci, A. Magnanini and A. Lo Nostro.** 2014. Prevalence and antibiotic resistance of *Enterococcus* spp. isolated from retail cheese, ready-to-eat salads, ham, and raw meat. *Food Microbiol.* 41: 1–7.
- Peters R.P.H., M.A. van Agtmael, S. Gierveld, S.A. Danner, A.B.J. Groeneveld, C.M.J.E. Vandenbroucke-Grauls and P.H.M. Savelkoul.** 2007. Quantitative detection of *Staphylococcus aureus* and *Enterococcus faecalis* DNA in blood to diagnose bacteremia in patients in the intensive care unit. *J. Clin. Microbiol.* 45(11): 3641–3646.
- Rehaïem A., Z. Ben Belgacem, M.R. Edalatian, B. Martínez, A. Rodríguez, M. Maná and N.P. Guerra.** 2014. Assessment of potential probiotic properties and multiple bacteriocin encoding-genes of the technological performing strain *Enterococcus faecium* MMRA. *Food Control* 37: 343–350.
- Reviriego C., T. Eaton, R. Martín, E. Jiménez, L. Fernández, M.J. Gasson and J.M. Rodríguez.** 2005. Screening of virulence determinants in *Enterococcus faecium* strains isolated from breast milk. *J. Hum. Lac.* 21(2): 131–137.
- Riboldi G.P., J. Frazzon, P.A. Azevedo and A.P.G. Frazzon.** 2009. Antimicrobial resistance profile of *Enterococcus* spp. isolated from food in southern Brazil. *Braz. J. Microbiol.* 40: 125–128.
- Rivas F.P., M.P. Castro, M. Vallejo, E. Marguet and C.A. Campos.** 2012. Antibacterial potential of *Enterococcus faecium* strains isolated from ewes' milk and cheese. *LWT – Food Sci. Technol.* 46: 428–436.
- Semedo T., M. Almeida, M.F. Silva Lopes, J.J. Figueiredo Marques, M.T. Barreto Crespo and R. Tenreiro.** 2003. Virulence factors in food, clinical and reference enterococci: a common trait in the genus? *Sys. Appl. Microbiol.* 26: 13–22.
- Singh K.V., S.R. Nallapareddy, J. Sillanpää and B.E. Murray.** 2010. Importance of the collagen adhesin Ace in pathogenesis and protection against *Enterococcus faecalis* experimental endocarditis. *PLoS Pathogens* 6: 1–13.
- Sparo M., L. Urbizu, M.V. Solana, G. Pourcel, G. Delpéch, A. Confalonieri, M. Ceci and S.F. Sanchez Bruni.** 2011. High-level resistance to gentamicin: genetic transfer between *Enterococcus faecalis* isolated from food of animal origin and human microbiota. *Let. Appl. Microbiol.* 54: 119–125.
- Theppangna W., T. Murase, N. Tokumaru, H. Chikumi, E. Shimizu and K. Otsuki.** 2007. Screening of the enterocin genes and antimicrobial activity against pathogenic bacteria in *Enterococcus* strains obtained from different origins. *J. Vet. Med. Sci.* 69(12): 1235–1239.
- Tuncer Y.** 2009. Some technological properties of phenotypically identified Enterococci Strains isolated from Turkish Tulum cheese. *Afr. J. Biotechnol.* 8(24): 7008–7016.
- Tuncer M., B. Özden Tuncer and Y. Tuncer.** 2014. Çiğ süttten izole edilen enterosin B üreticisi *Enterococcus faecalis* MYE58 suşunun güvenlik değerlendirmesi. *Gıda/The Journal of Food* 39(5): 275–282.
- Valenzuela A.S., N.B. Omar, H. Abriouel, R.L. López, E. Ortega, M.M. Cañamero and A. Gálvez.** 2008. Risk factors in enterococci isolated from foods in Morocco: Determination of antimicrobial resistance and incidence of virulence traits. *Food Chem. Toxic.* 46: 2648–2652.
- van Belkum M.J., B.J. Hayema, A. Geis, J. Kok and G. Venema.** 1989. Cloning of two bacteriocin genes from a lactococcal bacteriocin plasmid. *Appl. Environ. Microbiol.* 55: 1187–1191.
- Vankerckhoven V., T. Van Autgaerden, C. Vael, C. Lammens, S. Chapelle, R. Rossi, D. Jabes and H. Goossens.** 2004. Development of multiplex PCR for the detection of *asa1*, *gelE*, *cylA*, *esp*, and *hyl* genes in Enterococci and survey for virulence determinants among European hospital isolates of *Enterococcus faecium*. *J. Clin. Microbiol.* 42: 4473–4479.
- Yogurtcu N.N and Y. Tuncer.** 2013. Antibiotic susceptibility patterns of *Enterococcus* strains isolated from Turkish Tulum cheese. *Int. J. Dairy Technol.* 66(2): 236–242.
- Yousif N.M.K., P. Dawyndt, H. Abriouel, A. Wijaya, U. Schillinger, M. Vancanneyt, J. Swings, H.A. Dirar, W.H. Holzapfel and C.M.A.P. Franz.** 2005. Molecular characterization, technological properties and safety aspects of enterococci from “Hussuwa”, an African fermented sorghum product. *J. Appl. Microbiol.* 96: 216–228.
- Yüceer Ö. and B. Özden Tuncer.** 2015. Determination of antibiotic resistance and biogenic amine production of lactic acid bacteria isolated from fermented Turkish sausage (sucuk). *J. Food Safety* 35: 276–285.

