

Rapid Detection and Differentiation of KPC and MBL Carbapenemases among Enterobacterales Isolates by a Modified Combined-Disk Test


MING WEI[#], PENG WANG[#], SHUAI WANG, CHUNXIA YANG and LI GU^{*}

Department of Infectious Diseases and Clinical Microbiology, Beijing Chao-Yang Hospital, Capital Medical University, Beijing, People's Republic of China

Submitted 13 May 2021, revised 12 July 2021, accepted 28 July 2021

Abstract

This study was conducted to develop a cheap, rapid, and accurate modified combined-disk test (mCDT) approach to detect and differentiate KPC and MBL carbapenemases among clinical carbapenem-resistant Enterobacterales (CRE) isolates and simultaneously distinguish them from carbapenem-susceptible Enterobacterales (CSE) isolates. A total of 163 CRE and 90 third-generation cephalosporin-resistant Enterobacterales isolates were tested using imipenem and meropenem disks and different concentrations of carbapenemase inhibitors. The optimal sensitivity and specificity for detecting KPC carbapenemase were 97.2% and 100%, respectively. The sensitivity and specificity for detecting MBL carbapenemase were 100% and 100% with imipenem or meropenem and carbapenemase inhibitors within six hours. The inhibitory zone diameter of 18 mm for imipenem or meropenem disks without inhibitor could distinguish CRE from CSE isolates. Therefore, this mCDT approach may be a useful tool in clinical laboratories to



detect CRE isolates and differentiate KPC and MBL producers, which is beneficial for patient management and hospital infection prevention and control.

Key words: carbapenem-resistant Enterobacterales, carbapenemase, rapid detection and differentiation, modified combined-disk test

Introduction

Infections caused by carbapenem-resistant Enterobacterales (CRE) are a major clinical challenge and a public health problem (WHO 2017). According to the 2019 report of the Centers for Disease Control and Prevention (CDC), CRE can cause 13,100 infections and 1,100 deaths per year in the USA (CDC 2019). The resistance to carbapenems in Enterobacterales is mainly associated with the production of different classes of carbapenemases (Nordmann et al. 2012).

Horizontal transfer of carbapenemase coding genes through mobile genetic elements such as plasmids and transposons between Gram-negative microorganisms is responsible for the rapid increase of carbapenemase-producing CRE (CP-CRE) isolates. Carbapenemases

can hydrolyze not only carbapenems but also most other β -lactam antibiotics. There is often coexistence of additional resistance mechanisms to other antibiotic classes (e.g., fluoroquinolones and aminoglycosides) in CP-CRE isolates (van Duin and Doi 2017). It leads to very few treatment options available against these organisms (Nordmann et al. 2012; van Duin and Doi 2017). However, a few newly commercialized antibiotics (e.g., ceftazidime/avibactam and meropenem/vaborbactam) have been used for the treatment of *Klebsiella pneumoniae* carbapenemase (KPC) producers recently, but failed to treat metallo- β -lactamase (MBL) producers (King et al. 2017; Bassetti et al. 2018; Pfaller et al. 2018). Therefore, it is crucial to detect and differentiate KPC and MBL carbapenemase-producers rapidly in individual patients.

[#] Ming Wei and Peng Wang contributed equally to this study.

^{*} Corresponding author: L. Gu, Department of Infectious Diseases and Clinical Microbiology, Beijing Chao-Yang Hospital, Capital Medical University, Beijing, People's Republic of China; e-mail: guli2013227@foxmail.com

© 2021 Ming Wei et al.

This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Several carbapenemase detection methods have been developed, including carbapenem hydrolysis derived methods, such as the modified Hodge test (Carvalhoes et al. 2010), modified carbapenem inactivation method (mCIM) (Pierce et al. 2017), Carba NP test (Laolerd et al. 2018), matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry assay (Papagiannitsis et al. 2015), carbapenemase activity inhibition based methods (Li et al. 2019), and antibody- (Kieffer et al. 2019) and PCR-based methods (Cointe et al. 2019). However, modified Hodge test, mCIM, Carba NP test, and MALDI-TOF assays cannot differentiate KPC and MBL carbapenemase, although the latter two methods require only about two hours to diagnose (Papagiannitsis et al. 2015; Pierce et al. 2017; Laolerd et al. 2018). Although antibody- and PCR-based methods can differentiate KPC and MBL within two hours, they cannot distinguish CRE from carbapenem-susceptible Enterobacterales (CSE) isolates when encountering an Enterobacterales isolate whose antimicrobial susceptibility is unknown. Besides, they are more expensive than other methods (Cointe et al. 2019; Kieffer et al. 2019). As a result, all of these available methods cannot meet the clinical requirements satisfactorily.

KPC and MBL are the most frequently encountered carbapenemases among CP-CRE isolates in China, and the latter is distributed worldwide (Logan and Weinstein 2017). Therefore, in the context, we explored a cheap, rapid, and accurate method to detect and characterize KPC and MBL carbapenemases among Enterobacterales isolates and to discriminate CRE from CSE for providing the basis of choice of antibiotics for clinicians to treat CP-CRE infected patients and prevent their further spread in medical institutions.

Experimental

Materials and Methods

Bacterial isolates. A total of 253 retrospectively collected between January 2014 and January 2019, non-duplicate Enterobacterales isolates, including 163 CRE isolates and 90 third-generation cephalosporin-resistant Enterobacterales (3GCeRE) isolates from the Department of Infectious Diseases and Clinical Microbiology, Beijing Chao-Yang Hospital were included. CRE was defined as an isolate non-susceptible to imipenem or meropenem (for the bacteria intrinsically resistant to imipenem, non-susceptible to meropenem other than imipenem is required), with a minimum inhibitory concentration (MIC) ≥ 2 $\mu\text{g/ml}$, or producing carbapenemase. 3GCeRE was defined as an isolate resistant to ceftazidime (MIC ≥ 16 $\mu\text{g/ml}$), ceftriaxone

(MIC ≥ 4 $\mu\text{g/ml}$), and cefotaxime (MIC ≥ 4 $\mu\text{g/ml}$), but susceptible to carbapenems. The MICs were measured by the broth microdilution (BMD) method, and the interpretative criteria were based on the Clinical and Laboratory Standards Institute (CLSI 2019). All the isolates were identified by the Vitek MALDI-TOF MS (bioMérieux, France).

Molecular detection of carbapenemase genes.

As previously described, the *bla*_{IMP}, *bla*_{NDM} and *bla*_{VIM} genes were each detected by PCR (Jing et al. 2018). The *bla*_{NMC}, *bla*_{SME}, *bla*_{IMP}, *bla*_{GES}, *bla*_{SPM}, *bla*_{GIM}, *bla*_{SIM} and *bla*_{OXA-48-like} genes were detected by a single primer set (Queenan and Bush 2007). The primers for the *bla*_{KPC} gene used in the study were previously described (Poirel et al. 2011), but the PCR was performed by a different procedure. Briefly, 12.5 μl of PCR Master Mix (Thermo Scientific, USA) was mixed with 2 μl of forward and reverse primers and water to a final volume of 23 μl . Then, 2 μl of purified DNA template was added to the mix. The PCR program consisted of an initial denaturation step at 95°C for 5 min, followed by 35 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, elongation at 72°C for 60 s, and a final extension at 72°C for 7 min. All of the primers used in the study are listed in Table S1.

The PCR products were sequenced bi-directionally using an ABI 3730XL DNA sequencer (Applied Biosystems, USA). The gene sequences were compared with those in the database located at the NCBI blast server (<http://blast.ncbi.nlm.nih.gov>). A minimum of 99% sequence identity and 99% coverage threshold was deemed to confirm each gene.

Phenotypic detection of KPC and MBL by a modified combined disk test. Inhibitor solutions of 50 mg/ml APB (3-aminophenyl boronic acid hydrochloride, Sigma-Aldrich, USA), and 0.5 M EDTA (ethylenediaminetetraacetic acid disodium salt dihydrate, Sigma-Aldrich, USA) were filtered using a 0.22 μm filter membrane (Millipore, Germany), and stored at 4°C (Petropoulou et al. 2006; Doi et al. 2008). The modified combined disk test (mCDT) was carried out using four 10- μg imipenem (or meropenem) disks (Oxoid, UK), including a disk alone, a disk plus 5 μl (or 10 μl) of APB for KPC inhibition, a disk plus 5 μl (or 10 μl) of EDTA for MBL inhibition, and a disk plus both APB and EDTA for simultaneous inhibition of KPC and MBL (Tsakris et al. 2010; Pournaras et al. 2013). The four disks were placed onto Mueller-Hinton agar (Becton, Dickinson and Company, USA) plates inoculated with bacterial suspensions of 2.80 ± 0.15 McFarland optical density. The inhibition zones were measured after incubation for 6 hours at $35 \pm 2^\circ\text{C}$ in ambient air. An increase of ≥ 5 mm in the inhibition zone diameter of the imipenem (or meropenem) disk containing inhibitors (APB, EDTA, or both) in comparison to the same disk without the

Table I
Species distribution of different carbapenemase types among non-CP-CRE and 3GCeRE isolates.

Category (n)	Species							
	Kpn	Eco	Ecl	Kae	Cfr	Pre	Pmi	Sma
KPC (107)	104	1	0	1	1	0	0	0
MBL (36)								
NDM (30)	7	15	1	0	1	4	1	1
IMP (6)	3	0	3	0	0	0	0	0
OXA-48-like (1)	1	0	0	0	0	0	0	0
KPC+NDM (1)	1	0	0	0	0	0	0	0
Non-CP-CRE (18)	8	5	2	3	0	0	0	0
3GCeRE (90)	17	47	7	10	2	1	4	2
Total (253)	141	68	13	14	4	5	5	3

Kpn – *Klebsiella pneumoniae*, Eco – *Escherichia coli*, Ecl – *Enterobacter cloacae*, Kae – *Klebsiella aerogenes*, Cfr – *Citrobacter freundii*, Pre – *Providencia rettgeri*, Pmi – *Proteus mirabilis*, Sma – *Serratia marcescens*

corresponding inhibitor was suggestive of KPC, MBL, or both carbapenemases production, respectively. Quality control strains included *Escherichia coli* ATCC 25922, *Klebsiella pneumoniae* ATCC BAA-1705, and *Klebsiella pneumoniae* ATCC BAA-1706.

Statistical analysis. The sensitivity and specificity were determined to assess the performance of mCDT for the identification and differentiation of carbapenemase using PCR results as a standard. Data were analyzed using the VassarStats online software (VassarStats.net).

Results

Species distribution and carbapenemase genes.

One hundred and sixty-three CRE and ninety 3GCeRE isolates comprised eight bacterial species: *K. pneumo-*

niae (n = 141), *E. coli* (68), *Klebsiella aerogenes* (n = 14), *Enterobacter cloacae* (n = 13), *Proteus mirabilis* (n = 5), *Providencia rettgeri* (n = 5), *Citrobacter freundii* (n = 4), and *Serratia marcescens* (n = 3). The PCR and sequencing results are shown in Table I. One hundred and forty-five CRE isolates were confirmed to be carrying carbapenemase genes. Of them, the bla_{KPC} gene was the most often discovered carbapenemase gene (n = 107), followed by the bla_{NDM} gene in 30 isolates, bla_{IMP} in six isolates, and $bla_{OXA-48-like}$ in one isolate, as well as both bla_{KPC} and bla_{NDM} in one isolate. The remaining 18 CRE isolates and all 3GCeRE isolates were negative for carbapenemase genes.

Differentiation of CRE and 3GCeRE isolates. The distribution of inhibitory zone diameter (IZD) of the imipenem disk alone after six hours of incubation is shown in Fig. 1a. The IZDs of all CRE isolates were




Fig. 1. The inhibitory zone diameter distribution of a) imipenem, b) meropenem for Enterobacteriales isolates tested.




Fig. 2. Flow diagram of the mCDT for distinguishing CRE from CSE isolates, and differentiating KPC and/or MBL producers.

A – imipenem disk alone, B – imipenem disk plus 5 μ l EDTA, C – imipenem disk plus 10 μ l APB and D – imipenem disk plus 5 μ l EDTA and 10 μ l APB.

≤ 18 mm, while all 3GCeRE isolates were ≥ 19 mm. For three *P. mirabilis* the value of IZD was the same and equal to 18 mm. Therefore, apart from the three *P. mirabilis* isolates, the imipenem disk alone could distinguish CRE from all the other 3GCeRE isolates after six hours of incubation.

Compared with imipenem, the distribution of IZDs of meropenem disk alone provided a clear distinction of CRE and 3GCeRE isolates (Fig. 1b). The IZDs of all CRE isolates were ≤ 18 mm, while all 3GCeRE isolates were ≥ 20 mm.

Sensitivity and specificity of mCDT. The interpretation of the mCDT results is shown in Fig. 2. When the IZD value of imipenem or meropenem disk alone was

≥ 19 mm, the Enterobacteriales isolate was deemed CSE, otherwise deemed CRE. Subsequently, an increase of ≥ 5 mm in the IZD of the imipenem (or meropenem) disk containing inhibitors (APB, EDTA, or both) compared to the disk without the corresponding inhibitor is deemed as KPC among CRE isolates, MBL, or both carbapenemases producer, respectively. At the same time, an increase of < 5 mm was deemed as OXA-48-like carbapenemase or non-CP-CRE isolate. Fig. 3 shows examples of mCDT results.

The accuracies of the mCDT among CRE isolates are shown in Table II. For identification of KPC carbapenemase, the sensitivity of the IPM (imipenem)-5 μ l APB was 88.8% (95/107), with a specificity of 100%

Table II
Accuracy of detecting KPC or MBL carbapenemases among 163 CRE isolates under different conditions.

Category	Carbapenemase	Accuracy	
		Sensitivity (%), (95% CI)	Specificity (%), (95% CI)
IPM-5 μ l APB	KPC	88.8 (80.9–93.8)	100 (92.0–100)
IPM-5 μ l EDTA	MBL	100 (88.0–100)	100 (96.3–100)
IPM-10 μ l APB	KPC	97.2 (91.4–99.3)	100 (92.0–100)
IPM-10 μ l EDTA	MBL	100 (88.0–100)	100 (96.3–100)
MEM-5 μ l APB	KPC	48.6 (38.9–58.4)	100 (92.0–100)
MEM-5 μ l EDTA	MBL	100 (88.0–100)	100 (96.3–100)
MEM-10 μ l APB	KPC	59.8 (49.9–69.0)	100 (92.0–100)
MEM-10 μ l EDTA	MBL	100 (88.0–100)	100 (96.3–100)

IPM – imipenem, MEM – meropenem, APB – 3-aminophenyl boronic acid hydrochloride, EDTA – ethylenediaminetetraacetic acid disodium salt dihydrate




Fig. 3. Examples of mCDT showing the inhibitory zone diameters of A, B, C, and D for a) a plate 7, 7, 14, and 15 mm confirming the KPC-producing isolate, and for b) a plate 9, 17, 8, 18 mm confirming the MBL-producing isolate.

A – imipenem disk alone, B – imipenem disk plus 5 μ l EDTA, C – imipenem disk plus 10 μ l APB and D – imipenem disk plus 5 μ l EDTA and 10 μ l APB.

(56/56). One *C. freundii* and 11 *K. pneumoniae* isolates were a false negative. Three *K. pneumoniae* isolates were incorrectly identified as non-CP-CRE or OXA-48-like producers, while the remaining eight isolates were incorrectly identified as KPC and MBL producers. The sensitivity of the IPM-10 μ l APB was 97.2% (104/107), with a specificity of 100% (56/56). Three *K. pneumoniae* isolates were incorrectly identified as both KPC and MBL producers. These three organisms gave similar results with IPM-5 μ l APB group. However, the sensitivities of the MEM (meropenem)-5 μ l APB and MEM-10 μ l APB were 48.6% (52/107) and 59.8% (64/107), respectively, with specificities of 100% (56/56) in both groups. Tiny bacterial colonies were observed for many KPC-producing isolates around MEM disk with APB. Therefore, the sensitivities for MEM-5 μ l APB and MEM-10 μ l APB groups were much lower than that of IPM with APB groups. Fifty and 36 isolates were incorrectly identified as non-CP-CRE or OXA-48-like producers, while five and seven isolates were incorrectly identified as both KPC and MBL producers in MEM-5 μ l APB and MEM-10 μ l APB group respectively.

For the identification of MBL carbapenemases, the sensitivity of the IPM-5 μ l EDTA was 100% (36/36), with a specificity of 100% (127/127). The same results were obtained for IPM-10 μ l EDTA, MEM-5 μ l EDTA, and MEM-10 μ l EDTA groups.

No statistical analysis was performed for this type because there was only one isolate that produced both KPC and NDM carbapenemases. The isolate was correctly identified for all IPM or MEM disks with APB + EDTA groups.

Discussion

The current study detected five different genes by PCR and sequencing method in 88.9% of CRE isolates ($n = 163$). The results showed that the bla_{KPC} (73.8%) and bla_{NDM} (20.7%) genes were most common, followed by bla_{IMP} (4.1%), $bla_{OXA-48-like}$ (0.7%), and $bla_{KPC+NDM}$ (0.7%). It follows a report of nationwide surveillance of clinical CRE isolates in China, which showed 93% of clinical isolates ($n = 1105$) producing carbapenemases, with a majority of isolates producing KPC (57%) or NDM (31%) carbapenemases (Zhang et al. 2017). Therefore, the main carbapenem-resistant mechanism for clinical Enterobacterales isolates is acquiring the bla_{KPC} or bla_{NDM} genes in China.

Due to the high morbidity and mortality associated with the invasive infection of CP-CRE strains (CDC 2019), accurate and fast detection and differentiation of CP-CRE types are critical in the individual patient. It will help the physicians make decisions on appropriate antibiotic treatment and improve prevention and control, especially in outbreak situations (Gutiérrez-Gutiérrez et al. 2017; Livermore et al. 2018; Li et al. 2019). Although several methods have been proposed (Burckhardt and Zimmermann 2011; Findlay et al. 2015; Bogaerts et al. 2020) they cannot fully meet the clinical demands. The mCIM plus EDTA-carbapenem inactivation method (eCIM) and CDT require more than 20 hours. Besides, mCIM/eCIM could not detect MBL carbapenemase when an Enterobacterales isolate produced two types of carbapenemases, including the MBL carbapenemase (Tsai et al. 2020). While Carba NP and MALDI-TOF MS are rapid, it cannot

differentiate the type of a carbapenemase. In addition, commercial immunochromatographic assay and PCR method require high cost and cannot distinguish CRE from CSE.

In recent years, a simplified carbapenem inactivation method (sCIM) for detecting carbapenemases with high accuracy has been established (Jing et al. 2018; Yamada et al. 2021). Compared to mCIM, it is easier to carry out by smearing the test strain onto an imipenem/meropenem disk and then placed the disk on Mueller-Hinton agar streaked with 0.5 McFarland standard *Escherichia coli* ATCC 25922. The result was read after overnight incubation. However, similar to mCIM, the long incubation time and inability to distinguish the type of carbapenemases will limit its widespread use in clinical practice.

This study developed the mCDT method because APB can inhibit the KPC carbapenemase and EDTA can inhibit MBL carbapenemases (Petropoulou et al. 2006; Doi et al. 2008). A higher initial concentration of the bacteria tested will reduce the time to form visible bacterial colonies. However, when the initial bacterial suspension was set for 3.5 McFarland optical density, the inhibitory effect of APB on the KPC carbapenemase was significantly reduced for many isolates, and tiny bacterial colonies could be seen around an imipenem disk with APB (Fig. S1). Thus, the inoculum was 2.80 ± 0.15 McFarland optical density (less than 3) to prevent the tiny bacterial colonies in this study. The incubation time was set for six hours, making it easy to read results due to the formation of a clear inhibition zone compared to five hours or less. Therefore, it is a facile and inexpensive in-house approach, which requires only six hours because the bacteria's high concentration (2.80 ± 0.15 McFarland optical density) is inoculated.

The mCDT can be utilized to detect CRE in clinical laboratories and simultaneously differentiate the KPC and MBL type of CP-CRE isolates. An overnight pure culture of bacteria identified as Enterobacterales strains can be used for mCDT in the morning, and the result can be acquired after six hours of incubation in the afternoon. As a result, this method can shorten the turnaround time since it can be carried out simultaneously as the antibiotic susceptibility test and acquire the results in six hours. It should be noted that an IZD of 18 mm as the breakpoint we describe is smaller than the IZD of 22 mm for imipenem and meropenem proposed by CLSI to distinguishing CRE from CSE isolates (CLSI 2019). In this study, applying the 18 mm breakpoint could detect all CRE strains and exclude all 3GCeRE strains for meropenem disk, but the IZDs of imipenem disk for three carbapenem-susceptible *P. mirabilis* isolates were 18 mm due to the intrinsic imipenem resistance. It is observed for *Morganella morganii*, *Proteus* spp., and *Providencia* spp. (CDC 2019).

Therefore, it is better to use a meropenem disk to distinguish CRE from CSE isolates for those intrinsically imipenem-resistant strains.

APB inhibits KPC carbapenemase due to the boronate moiety binding to the catalytic serine side chain; thus, forming the covalent adduct and inactivating it (Hecker et al. 2015). In the current study, most KPC-producing isolates could not be detected by meropenem but imipenem disk due to tiny bacterial colonies formed around a meropenem disk with APB. This result indicates that the covalent adduct still retains a low level of catalytic efficiency, and the catalytic efficiency to meropenem is higher than that of imipenem. The reason may be that the affinity of the covalent adduct to meropenem is higher than that of imipenem.

This study has some limitations. First, the mCDT was not suitable for detecting OXA-48-like carbapenemases because there is no specific inhibitor against OXA-48-like carbapenemases available. Second, this study did not include MBL carbapenemases except NDM and IMP (e.g. VIM and others). Third, this method cannot detect class A carbapenemases except KPC (e.g. GES, SME) because they cannot be inhibited by APB (Nordmann et al. 2012). According to the report of Lee and Suh (2021), detection of GES by modified Hodge test (MHT), mCIM, and Rapidec Carba NP was also very poor, which is a tricky problem.

Conclusions

In summary, the proposed mCDT that used 10 μ l APB and 5 μ l EDTA as inhibitors enables detection and differentiation of KPC and MBL carbapenemase types and distinguishes CRE from CSE simultaneously within six hours among Enterobacterales isolates. It could be a low cost, high accuracy, and easy operational approach. While this approach requires validation in other laboratories, we believe that the mCDT may be useful, especially in low-income countries and regions where KPC and MBL carbapenemases are epidemics.

Acknowledgments

The authors would like to thank all staff members of the Department of Infectious Diseases and Clinical Microbiology, Beijing Chao-Yang Hospital (Beijing, China), for their contribution to this work.

Author contributions

LG and MW conceived the study. MW, PW, SW and CY performed the research. LG, MW, and PW analyzed data and wrote the paper.

Funding

This work was supported by the Capital Health Research and Development of Special (Grant number 2018-1-4081).

Conflict of interest

The authors do not report any financial or personal connections with other persons or organizations, which might negatively affect the contents of this publication and/or claim authorship rights to this publication.

Literature

- Bassetti M, Giacobbe DR, Giamarellou H, Viscoli C, Daikos GL, Dimopoulos G, De Rosa FG, Giamarellos-Bourboulis EJ, Rossolini GM, Righi E, et al.; Critically Ill Patients Study Group of the European Society of Clinical Microbiology and Infectious Disease (ESCMID); Hellenic Society of Chemotherapy (HSC); Società Italiana di Terapia Antinfettiva (SITA). Management of KPC-producing *Klebsiella pneumoniae* infections. *Clin Microbiol Infect*. 2018 Feb;24(2):133–144. <https://doi.org/10.1016/j.cmi.2017.08.030>
- Bogaerts P, Berger AS, Evrard S, Huang TD. Comparison of two multiplex immunochromatographic assays for the rapid detection of major carbapenemases in Enterobacterales. *J Antimicrob Chemother*. 2020 Jun 01;75(6):1491–1494. <https://doi.org/10.1093/jac/dkaa043>
- Burckhardt I, Zimmermann S. Using matrix-assisted laser desorption ionization-time of flight mass spectrometry to detect carbapenem resistance within 1 to 2.5 hours. *J Clin Microbiol*. 2011 Sep; 49(9):3321–3324. <https://doi.org/10.1128/JCM.00287-11>
- Carvalhoes CG, Pição RC, Nicoletti AG, Xavier DE, Gales AC. Cloverleaf test (modified Hodge test) for detecting carbapenemase production in *Klebsiella pneumoniae*: be aware of false positive results. *J Antimicrob Chemother*. 2010 Feb 01;65(2):249–251. <https://doi.org/10.1093/jac/dkp431>
- CDC. Antibiotic resistance threats in the United States, 2019. Atlanta (USA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2019. <https://doi.org/10.15620/cdc:82532>
- CLSI. M100-S29 Performance standards for antimicrobial susceptibility testing. Wayne (USA): Clinical and Laboratory Standards Institute; 2019.
- Cointe A, Walewski V, Hobson CA, Doit C, Bidet P, Dortet L, Bonacorsi S, Birgy A. Rapid carbapenemase detection with Xpert Carba-R V2 directly on positive blood vials. *Infect Drug Resist*. 2019 Oct;12:3311–3316. <https://doi.org/10.2147/IDR.S204436>
- Doi Y, Potoski BA, Adams-Haduch JM, Sidjabat HE, Pasculle AW, Paterson DL. Simple disk-based method for detection of *Klebsiella pneumoniae* carbapenemase-type β -lactamase by use of a boronic acid compound. *J Clin Microbiol*. 2008 Dec;46(12):4083–4086. <https://doi.org/10.1128/JCM.01408-08>
- Findlay J, Hopkins KL, Meunier D, Woodford N. Evaluation of three commercial assays for rapid detection of genes encoding clinically relevant carbapenemases in cultured bacteria. *J Antimicrob Chemother*. 2015 May;70(5):1338–1342. <https://doi.org/10.1093/jac/dku571>
- Gutiérrez-Gutiérrez B, Salamanca E, de Cueto M, Hsueh PR, Viale P, Paño-Pardo JR, Venditti M, Tumbarello M, Daikos G, Cantón R, et al.; REIPI/ESGBIS/INCREMENT Investigators. Effect of appropriate combination therapy on mortality of patients with bloodstream infections due to carbapenemase-producing Enterobacteriaceae (INCREMENT): a retrospective cohort study. *Lancet Infect Dis*. 2017 Jul;17(7):726–734. [https://doi.org/10.1016/S1473-3099\(17\)30228-1](https://doi.org/10.1016/S1473-3099(17)30228-1)
- Hecker SJ, Reddy KR, Totrov M, Hirst GC, Lomovskaya O, Griffith DC, King P, Tsvikovski R, Sun D, Sabet M, et al. Discovery of a cyclic boronic acid β -lactamase inhibitor (RPX7009) with utility vs class a serine carbapenemases. *J Med Chem*. 2015 May 14;58(9): 3682–3692. <https://doi.org/10.1021/acs.jmedchem.5b00127>
- Jing X, Zhou H, Min X, Zhang X, Yang Q, Du S, Li Y, Yu F, Jia M, Zhan Y, et al. The simplified carbapenem inactivation method (sCIM) for simple and accurate detection of carbapenemase-producing gram-negative bacilli. *Front Microbiol*. 2018 Oct 30;9:2391. <https://doi.org/10.3389/fmicb.2018.02391>
- Kieffer N, Poirel L, Nordmann P. Rapid immunochromatography-based detection of carbapenemase producers. *Infection*. 2019 Aug; 47(4):673–675. <https://doi.org/10.1007/s15010-019-01326-1>
- King M, Heil E, Kuriakose S, Bias T, Huang V, El-Beyrouty C, McCoy D, Hiles J, Richards L, Gardner J, et al. Multicenter study of outcomes with ceftazidime-avibactam in patients with carbapenem-resistant Enterobacteriaceae infections. *Antimicrob Agents Chemother*. 2017 Jul;61(7):e00449–e17. <https://doi.org/10.1128/AAC.00449-17>
- Laolerd W, Akeda Y, Preeyanon L, Rattawongjirakul P, Santanirand P. Carbapenemase-producing carbapenem-resistant Enterobacteriaceae from Bangkok, Thailand, and their detection by the Carba NP and modified carbapenem inactivation method tests. *Microb Drug Resist*. 2018 Sep;24(7):1006–1011. <https://doi.org/10.1089/mdr.2018.0080>
- Lee AJ, Suh HS. Comparative evaluation of the modified carbapenem inactivation method for phenotypic detection of Guiana extended-spectrum β -lactamase-type carbapenemases in Enterobacterales. *Lab Med*. 2021 Apr 30;lmab026. <https://doi.org/10.1093/labmed/lmab026>
- Li J, Li C, Cai X, Shi J, Feng L, Tang K, Tong Y, Li Y. Performance of modified carbapenem inactivation method and inhibitor-based combined disk test in the detection and distinguishing of carbapenemase producing Enterobacteriaceae. *Ann Transl Med*. 2019 Oct; 7(20):566. <https://doi.org/10.21037/atm.2019.09.43>
- Livermore DM, Meunier D, Hopkins KL, Doumith M, Hill R, Pike R, Staves P, Woodford N. Activity of ceftazidime/avibactam against problem Enterobacteriaceae and *Pseudomonas aeruginosa* in the UK, 2015–16. *J Antimicrob Chemother*. 2018 Mar 01;73(3):648–657. <https://doi.org/10.1093/jac/dkx438>
- Logan LK, Weinstein RA. The epidemiology of carbapenem-resistant Enterobacteriaceae: the impact and evolution of a global menace. *J Infect Dis*. 2017 Feb 15;215(suppl_1):S28–S36. <https://doi.org/10.1093/infdis/jiw282>
- Nordmann P, Dortet L, Poirel L. Carbapenem resistance in Enterobacteriaceae: here is the storm! *Trends Mol Med*. 2012 May;18(5): 263–272. <https://doi.org/10.1016/j.molmed.2012.03.003>
- Papagiannitsis CC, Študentová V, Izdebski R, Oikonomou O, Pfeifer Y, Petinaki E, Hrabák J. Matrix-assisted laser desorption ionization-time of flight mass spectrometry meropenem hydrolysis assay with NH_4HCO_3 , a reliable tool for direct detection of carbapenemase activity. *J Clin Microbiol*. 2015 May;53(5):1731–1735. <https://doi.org/10.1128/JCM.03094-14>
- Petropoulou D, Tzanetou K, Syriopoulou VP, Daikos GL, Ganteris G, Malamou-Lada E. Evaluation of imipenem/imipenem + EDTA disk method for detection of metallo- β -lactamase-producing *Klebsiella pneumoniae* isolated from blood cultures. *Microb Drug Resist*. 2006 Mar;12(1):39–43. <https://doi.org/10.1089/mdr.2006.12.39>
- Pfaller MA, Huband MD, Mendes RE, Flamm RK, Castanheira M. *In vitro* activity of meropenem/vaborbactam and characterisation of carbapenem resistance mechanisms among carbapenem-resistant Enterobacteriaceae from the 2015 meropenem/vaborbactam surveillance programme. *Int J Antimicrob Agents*. 2018 Aug;52(2):144–150. <https://doi.org/10.1016/j.ijantimicag.2018.02.021>
- Pierce VM, Simner PJ, Lonsway DR, Roe-Carpenter DE, Johnson JK, Brasso WB, Bobenchik AM, Lockett ZC, Charnot-Katsikas A,

- Ferraro MJ, et al.** Modified carbapenem inactivation method for phenotypic detection of carbapenemase production among *Enterobacteriaceae*. *J Clin Microbiol*. 2017 Aug;55(8): 2321–2333. <https://doi.org/10.1128/JCM.00193-17>
- Poirel L, Walsh TR, Cuvillier V, Nordmann P.** Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis*. 2011 May;70(1):119–123. <https://doi.org/10.1016/j.diagmicrobio.2010.12.002>
- Pournaras S, Zarkotou O, Poulou A, Kristo I, Vrioni G, Themeli-Digalaki K, Tsakris A.** A combined disk test for direct differentiation of carbapenemase-producing *Enterobacteriaceae* in surveillance rectal swabs. *J Clin Microbiol*. 2013 Sep;51(9):2986–2990. <https://doi.org/10.1128/JCM.00901-13>
- Queenan AM, Bush K.** Carbapenemases: the versatile β -lactamases. *Clin Microbiol Rev*. 2007 Jul;20(3):440–458. <https://doi.org/10.1128/CMR.00001-07>
- Tsai YM, Wang S, Chiu HC, Kao CY, Wen LL.** Combination of modified carbapenem inactivation method (mCIM) and EDTA-CIM (eCIM) for phenotypic detection of carbapenemase-producing *Enterobacteriaceae*. *BMC Microbiol*. 2020 Dec;20(1):315. <https://doi.org/10.1186/s12866-020-02010-3>
- Tsakris A, Poulou A, Pournaras S, Voulgari E, Vrioni G, Themeli-Digalaki K, Petropoulou D, Sofianou D.** A simple phenotypic method for the differentiation of metallo- β -lactamases and class A KPC carbapenemases in *Enterobacteriaceae* clinical isolates. *J Antimicrob Chemother*. 2010 Aug 01;65(8):1664–1671. <https://doi.org/10.1093/jac/dkq210>
- van Duin D, Doi Y.** The global epidemiology of carbapenemase-producing *Enterobacteriaceae*. *Virulence*. 2017 May 19;8(4):460–469. <https://doi.org/10.1080/21505594.2016.1222343>
- WHO.** Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug resistant bacterial infections, including tuberculosis [Internet]. Geneva (Switzerland): World Health Organization; 2017 [cited 2021 Apr 14]. Available from <https://www.who.int/publications/i/item/WHO-EMP-IAU-2017.12>
- Yamada K, Sasaki M, Murakami H, Aoki K, Morita T, Ishii Y, Tateda K.** Evaluation of the simplified carbapenem inactivation method as a phenotypic detection method for carbapenemase-producing *Enterobacteriales*. *J Microbiol Methods*. 2021 Aug;187:106273. <https://doi.org/10.1016/j.mimet.2021.106273>
- Zhang R, Liu L, Zhou H, Chan EW, Li J, Fang Y, Li Y, Liao K, Chen S.** Nationwide surveillance of clinical carbapenem-resistant *Enterobacteriaceae* (CRE) strains in China. *EBioMedicine*. 2017 May; 19:98–106. <https://doi.org/10.1016/j.ebiom.2017.04.032>

Supplementary materials are available on the journal's website.